



Changes in Tree Species Abundance in a Neotropical Forest: Impact of Climate Change

Richard Condit, Stephen P. Hubbell, Robin B. Foster

Journal of Tropical Ecology, Volume 12, Issue 2 (Mar., 1996), 231-256.

Stable URL:

<http://links.jstor.org/sici?sici=0266-4674%28199603%2912%3A2%3C231%3ACITSAI%3E2.0.CO%3B2-0>

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

Journal of Tropical Ecology is published by Cambridge University Press. Please contact the publisher for further permissions regarding the use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/cup.html>.

Journal of Tropical Ecology
©1996 Cambridge University Press

JSTOR and the JSTOR logo are trademarks of JSTOR, and are Registered in the U.S. Patent and Trademark Office. For more information on JSTOR contact jstor-info@umich.edu.

©2002 JSTOR

Changes in tree species abundance in a Neotropical forest: impact of climate change

RICHARD CONDIT*, STEPHEN P. HUBBELL*† and ROBIN B. FOSTER*‡

* *Smithsonian Tropical Research Institute, Unit 0948, APO AA 34002-0948, USA; or Apartado 2072, Balboa, Ancón, Panamá*

† *Department of Ecology, Evolution and Behavior, Princeton University, Princeton, New Jersey 08544, USA*

‡ *Department of Botany, Field Museum of Natural History, Chicago, Illinois 60605, USA*

ABSTRACT. The abundance of all tree and shrub species has been monitored for eight years in a 50 ha census plot in tropical moist forest in central Panama. Here we examine population trends of the 219 most numerous species in the plot, assessing the impact of a long-term drying trend. Population change was calculated as the mean rate of increase (or decrease) over eight years, considering either all stems ≥ 10 mm diameter at breast height (dbh) or just stems ≥ 100 mm dbh. For stems ≥ 10 mm, 40% of the species had mean growth rates $< 1\%$ per year (either increasing or decreasing) and 12% had changes $\geq 5\%$ per year. For stems ≥ 100 mm, the figures were 38% and 8%.

Species that specialize on the slopes of the plot, a moist microhabitat relative to the plateau, suffered significantly more declines in abundance than species that did not prefer slopes (stems ≥ 10 mm dbh). This pattern was due entirely to species of small stature: 91% of treelets and shrubs that were slope-specialists declined in abundance, but just 19% of non-slope treelets and shrubs declined. Among larger trees, slope and non-slope species fared equally. For stems ≥ 100 mm dbh, the slope effect vanished because there were few shrubs and treelets with stems ≥ 100 mm dbh. Another edaphic guild of species, those occurring preferentially in a small swamp in the centre of the plot, were no more likely to decline in abundance than non-swamp species, regardless of growth form. Species that preferentially colonize canopy gaps in the plot were slightly more likely to decrease in abundance than non-colonizing species (only for stems ≥ 10 mm dbh, not ≥ 100 mm). Despite this overall trend, however, several colonizing species had the most rapidly increasing populations in the plot.

The impact of a 25-year drying trend and an associated increase in the severity of the 4-month dry season is having an obvious impact on the BCI forest. At least 16 species of shrubs and treelets with affinities for moist microhabitats are headed for extinction in the plot. Presumably, these species invaded the forest during a wetter period prior to 1966. A severe drought of 1983 that caused unusually high tree mortality contributed to this trend, and may also have been responsible for sharp increases in abundance of a few gap-colonizers because it temporarily opened the forest canopy. The BCI forest is remarkably sensitive to a subtle climatic shift, yet we do not know whether this is typical for tropical forests because no other large-scale censuses exist for comparison.

KEY WORDS: demography, population dynamics, tropical populations

INTRODUCTION

The myth that tropical climates provide a stable environment for tropical forest organisms has long been buried. We know that annual shifts in moisture availability can stress vegetation and limit the distribution of many species

(Hartshorn 1992, Wright 1992). Now, thanks to long-term weather records kept by the Panama Canal Commission, we also know that the vegetation of Barro Colorado Island (BCI) must face supra-annual shifts in moisture availability (Windsor 1990, Windsor *et al.* 1990). Total precipitation at BCI underwent an abrupt decline around 1965, averaging 2740 mm prior to 1965 and 2430 mm since, paralleling a worldwide reduction in rainfall in the northern tropics (Bradley *et al.* 1987, Diaz *et al.* 1989). Moreover, associated with a strong El Niño event, the annual dry season of 1983 was unusually long and severe, causing elevated tree mortality (Condit *et al.* 1992b, Condit *et al.* 1995, Leigh *et al.* 1990). The 1983 drought and the long-term drop in rainfall are undoubtedly part of the same phenomenon, as the frequency of dry seasons (15 December to 15 April) receiving less than 100 mm of rain increased from once every 6.2 years prior to 1965 to once every 3.5 years since (Windsor 1990).

How does the composition of a tropical forest change when precipitation patterns shift? Unusual droughts and variation in rainfall are recognized more and more as important in tropical forests (Foster 1982a, Hartshorn 1992, Leigh *et al.* 1990, Woods 1989), but we know little about how populations of individual species change as a result. Certainly we know that past climate changes have led to shifts in species' distributions (Bush & Colinvaux 1990, Bush *et al.* 1990, Hamilton & Taylor 1991, Sukumar *et al.* 1993), and in temperate forests, detailed descriptions of range shifts that accompany past climate changes are so well documented (Davis 1981, Delcourt & Delcourt, 1987) that precise predictions on the impact of future climate scenarios can be made (Botkin & Nisbet 1992, Dale & Franklin 1989, Franklin *et al.* 1992, Overpeck *et al.* 1990, Pastor & Post 1988, Shugart & Smith 1992, Solomon 1986, Urban *et al.* 1993). The species-specific information behind these predictions is not available for most tropical forests. Only for the Luquillo forest in Puerto Rico has a species-specific model been used to predict the impact of climate change; O'Brien *et al.* (1992) assessed the potential impact of increasing hurricane frequency.

To gather information on many individual species of tropical trees in one community, we established a large-scale and long-term population survey of forest at Barro Colorado Island in Panama. In order to include substantive information on populations of many species, a large plot was mapped: 50 ha of forest, with all stems above 10 mm in diameter included (Condit 1995, Hubbell & Foster 1983). This dataset provides detailed information on change in forest composition and its relation to climate change. Similar large plots in natural forest are now being censused in India, Malaysia, Thailand, Sri Lanka, Puerto Rico, Ecuador, Cameroon and Zaïre (Condit 1995, Manokaran *et al.* 1992, Sukumar *et al.* 1992, Zimmerman *et al.* 1994), so we will soon be able to make a worldwide assessment on the lability of the species composition of tropical forests.

Here we provide population estimates for 313 species of tropical trees found in the 50 ha plot in Panama between 1982 and 1990. We address specific hypotheses about how the community is changing, in particular, how it might be

affected by long-term reduction in rainfall (Condit *et al.* 1992b, Hubbell & Foster 1990a, 1992). First, we consider species whose distributions within the plot are associated with moist microhabitats: a seasonal swamp and the moderately sloping terrain that drops off from the plateau in the centre of the plot (Becker *et al.* 1988, Hubbell & Foster 1983, 1986a). These areas remain wet throughout most dry seasons because a basalt cap below the plateau accumulates water during the wet season and drains slowly into the swamp and slopes throughout the dry season. Our prediction is that species associated with the moist microsites should be especially sensitive to the overall drying trend and will have suffered disproportionate losses in population.

In addition, we consider population changes of species that preferentially colonize light gaps within the forest. There are two different predictions about colonizing species. First, Hubbell & Foster (1990a, 1992) suggested that the plot is undergoing a slow loss of weedy species, because the region just north of the plot (plus 2 ha within the plot) was cleared of forest about 90 years ago, and has since regrown. Colonizing species probably gained abundance within the old forest because of their large populations just outside, and are now declining. If this is the case, we should be able to detect disproportionate population declines among colonizing species. The second prediction on colonizers is just the opposite, and is based on the observation that the drought opened the forest canopy briefly during 1983 (Becker & Smith 1990). With more light reaching the ground, colonizing species should increase in abundance. Population changes of species preferring gaps can tell us which of the potentially opposing forces is more important.

MATERIALS AND METHODS

Study site

Barro Colorado (BCI) is a 1500 ha island that was a hilltop until the Panama Canal was finished in 1914. The island is part of the Barro Colorado Nature Monument and has been operated as a research reserve since 1923. It is entirely forested, most in old-growth forest with no signs of human disturbance for over 500 years: 48 ha of the 50 ha plot are in old-growth, with 2 ha in an area cleared until about 1900 as part of a French settlement. Temperatures are uniform year-round at BCI, but rainfall is seasonal, with almost none falling between mid-December and mid-April. Details on climate, flora and fauna can be found in Croat (1978) and Leigh *et al.* (1982).

Census

A 50 ha plot on the top of the island was fully censused in 1981–1983, 1985 and 1990 (Condit *et al.* 1992a,b, 1993a,b, Hubbell & Foster 1983, 1986a,b, 1987, 1990a,b, 1992); we refer to the first census, which lasted two years, as the 1982 census. All free-standing, woody stems ≥ 10 mm diameter at breast height (dbh) were identified, tagged and mapped. The diameter of each stem

was measured at breast height (1.3 m) unless there were irregularities in the trunk there, in which case the measurement was taken at the nearest point downward where the stem was cylindrical. Dbhs of buttressed trees were taken above the buttresses. There were about 242,000 living stems in each census (Hubbell & Foster 1990a), and 305,875 stems over all three censuses; 28 have not been identified to species. A total of 313 species have been identified: 304, 306 and 303 in successive censuses. (Three new species have been added since Condit *et al.* 1992b, all rare plants that had been misidentified as more common species.) Included in the list of 313 is a single tree that appeared to be a hybrid between *Apeiba membranacea* and *A. tibourbou*, and two distinct varieties of *Swartzia simplex* (Croat 1978).

Analyses

Species included. Abundances for all 313 species are reported. Species' names match those from Croat (1978) and D'Arcy (1987), except for species which were discovered, or whose names have been changed, since. An Appendix lists all cases where names do not match those found in Croat (1978) or D'Arcy (1987), and allows any species listed here to be located in those floras or in our previous publications on the 50 ha plot. Authorities for all species can be found via these references.

Tests of hypotheses about changes in abundance included only those species that had at least 20 individuals ≥ 10 mm dbh in at least one of the censuses. We used this cutoff because large percentage changes in abundance of very rare species could be caused by minor, chance events. Four species of *Bactris* palms were also eliminated from analyses because we changed methods for counting individuals of these species. This left 219 species for analyses of all stems ≥ 10 mm dbh. Analyses were then repeated for changes in the number of individuals ≥ 100 mm dbh, including the 136 species that had at least 20 stems in at least one census. We included an analysis with this larger cutoff because many other studies of tropical forest use the 100 mm limit (Phillips & Gentry 1994, Phillips *et al.* 1994).

Species characteristics. We analysed changes in abundance as they correlated with three species characteristics – growth form, moisture preference and tendency to recruit into light gaps. Species were divided into four growth forms – large trees (≥ 20 m tall), mid-sized trees (10–20 m), treelets (4–10 m) and shrubs (1–4 m) – based on the maximum height attained at BCI (Hubbell & Foster 1986a). Moisture regime was defined using the slopes in the 50 ha plot, which have higher soil moisture content during the dry season than the plateau above them (Becker *et al.* 1988), and the swamp, which is flooded throughout the wet season and remains moist in the dry season (Hubbell & Foster 1986a).

Many species have distributions clearly demarcated by the slopes and the swamp (Hubbell & Foster 1986a), and we calculated the density of all species in the different habitats (unpublished data). We used the ratio of density on

the slopes (all 20 m × 20 m quadrats inclined $\geq 7^\circ$) to density on the lower part of the plateau (quadrats with slope $< 7^\circ$ and elevation < 152 m, excluding the swamp) as an index of 'slope-specialization', and the ratio of density in the swamp (all 20 m × 20 m quadrats holding standing water through most of the wet season) to density on the lower plateau as an index for 'swamp-specialization'. We considered 'slope-specialists' and 'swamp-specialists' species with ratios ≥ 1.5 ; this cutoff was chosen because chi-squared tests showed that nearly all higher ratios were significantly different from 1.0 ($P < 0.01$), while most below did not (unpublished data). This index was preferable to a definition based on statistical significance, because the latter is sensitive to sample size. Finally, as a 'colonizing index' for each species, we used the fraction of recruits in light gaps given in Welden *et al.* (1991): Hubbell & Foster (1986b) used a similar but not identical 'index of heliophily'. Colonizers were defined as those species with an index ≥ 30 ; again, this corresponds roughly with a statistically significant preference for recruiting in gaps (Welden *et al.* 1991) but does not depend on sample size. Most colonizers are probably 'pioneers' as defined by Swaine & Whitmore (1988), but they emphasized seed germination characteristics, which we do not consider here. Species for which information was lacking were omitted from all analyses requiring that information. The slope and swamp indices were calculated for all but 27 of the very rare species, but the colonizing index was available for only the 156 species listed in Welden *et al.* (1991).

Statistical tests. In order to determine whether certain groups of species suffered disproportionate losses, the number of species that increased or decreased in abundance between 1982 and 1990 was tallied as a function of the four categorical variables. For statistical tests, a standard ANOVA was not possible because the design was unbalanced, with many empty cells. Instead, chi-squared tests were used on each of the variables: for example, a 2×2 contingency table for slope-specialization category and for population change provided a chi-squared statistic with one degree of freedom. To determine effects of each variable separately, we proceeded as follows. Swamp effect was assessed by contingency tables for swamp and non-swamp species; since swamp status was not associated with colonizing nor slope status, the latter two categories were simply ignored when swamp effect was tested. But slope and colonizing variables were associated – there were fewer slope-colonizer species than expected by chance – so we segregated species simultaneously by both categories. All tests were carried out on the four growth forms separately.

For each species, we calculated the annualized rate of population change (r) using a standard model of exponential population growth:

$$r = \frac{\ln N_t - \ln N_0}{t}$$

where N_t and N_0 are population sizes at time t and time 0 and \ln means the natural logarithm. The time interval, t , for each species was defined as the

arithmetic mean time elapsed between censuses for individuals of that species (based on the census data of each 20 m \times 20 m quadrat in the plot).

Earlier publications

Hubbell & Foster (1990a, 1992) described population changes based on the 1982–1985 interval, and Condit *et al.* (1992b) updated this with 1990 data, but this is the first presentation on abundance for all 313 species. Discrepancies between the numbers reported here and those from earlier reports are slight and are due solely to corrections of old errors. Since this is an on-going process, future reports might give figures slightly different from those reported here.

Access to data

We hope that the table of abundances for 313 species provided here will be useful for many future studies, and we will provide computer versions of the table to anyone interested. Please send us a diskette and indicate preferred formats.

RESULTS

Changes in abundance for stems ≥ 10 mm dbh

Of the 219 more common species considered here, 105 had increases in stem number between 1982 and 1990, 108 had decreases and six did not change. For all 313 species in the plot, 136 increased, 154 decreased and 23 did not change. As previously noted (Condit *et al.* 1992b, Hubbell & Foster 1990a, 1992), rare species – in this case the 94 having fewer than 20 stems in all censuses – suffered proportionally more declines than common species. Table 1 gives the abundance in all three censuses for all 313 species.

Many populations did not change by much (Figure 1). Of 219 species, 88 (40%) had population changes $< 1\%$ per year between 1982 and 1990. But some species had dramatic changes in abundance: 27 species (12%) changed at rates more than 5% per year, 18 declining and nine increasing (Table 2). Some common species underwent substantial declines. *Poulsenia armata* fell from 3430 to 2126 stems, and *Acalypha diversifolia* from 1568 to 827. The most rapid rate of decline was *Piper aequale*, which had 219 stems in 1982 and 83 in 1990 (Table 2). On the other hand, the population of *Palicourea guianensis* rose from 377 to 1475 stems, while the much less common *Psychotria graciflora* had the biggest rate of increase, from 10 to 44 stems over eight years (Table 2). The mean rate of change for the 219 species was -0.29% per year, while the mean rate of absolute change was 2.25% (the mean of the absolute values of rates of change).

Changes in abundance for stems ≥ 100 mm dbh

The range of population change among larger stems was no different than for smaller (Figure 1B). Of the 136 species considered, 66 increased in abundance from 1982 to 1990, 61 decreased and seven stayed the same. Fifty-one




Figure 1. Distribution of population growth rates. The 'colonizer' category includes all colonizing species that were not slope-specialists, and the 'slope' category includes slope species that were non-colonizing; 'slope-colonizers' are the four specializing in both areas. 'Other' includes all species that were neither slope nor colonizing plus all species that were missing information on one or both categories. (A) Rate of change of populations of stems ≥ 10 mm dbh, including 219 species (see text). (B) Rate of change of populations of stems ≥ 100 mm dbh, including 136 species (see text).

Table 1. Total populations of all 313 species recorded in the 50 ha plot on Barro Colorado Island, above two dbh cutoffs. Following each species name are four codes: a column headed G (for growth form), with T = large tree, M = mid-sized tree, U = shrub, H = hemi-epiphyte and ... = unknown; column headed C (colonizing status), W (swamp status) and S (slope status), under which, the letters C, W and S indicate specialists, species that were substantially more abundant in the given habitat (see text), the letter n indicates non-specialists, or species equally abundant inside or outside the given habitat (or more abundant outside), and ... indicates that information was not available (see text). The columns headed 'rate' give the exponential rate of population change between 1982 and 1990, first for stems ≥ 10 mm and then for stems ≥ 100 mm dbh.

Species	Codes				Family	Stems ≥ 10 mm dbh				Stems ≥ 100 mm dbh				Rate 1982-1990	Rate 1982-1990
	G	C	W	S		1982	1985	1990	Rate 1982-1990	1982	1985	1990	Rate 1982-1990		
<i>Acacia melanocera</i>	U	...	W	n	5	6	11	0.092	2	1	2	0.000			
<i>Acalypha diversifolia</i>	S	C	W	n	1568	1208	827	-0.072	2	3	3	0.044			
<i>Acalypha macrostachya</i>	U	C	n	n	80	67	45	-0.071	2	0	2	0.000			
<i>Adelia triloba</i>	U	C	n	n	345	314	279	-0.024	115	114	106	-0.009			
<i>Aegiphila panamensis</i>	M	n	n	n	134	125	93	-0.043	23	21	24	0.004			
<i>Alchornea costaricensis</i>	T	C	n	n	384	313	265	-0.043	160	158	157	-0.002			
<i>Alchornea latifolia</i>	M	...	n	n	2	2	3	0.052	2	2	2	0.000			
<i>Alibertia edulis</i>	U	n	W	n	305	342	377	0.024	2	2	3	0.046			
<i>Allophylus psilospermus</i>	M	n	n	n	175	171	153	-0.015	30	32	27	-0.012			
<i>Alseis blackiana</i>	T	n	n	n	7595	8057	8424	0.012	847	858	940	0.011			
<i>Amatoua corymbosa</i>	U	...	n	n	31	32	32	0.003	2	3	4	0.074			
<i>Anacardium excelsum</i>	T	...	W	S	28	26	24	-0.016	23	23	22	-0.004			
<i>Anaxagorea panamensis</i>	S	C	n	S	472	473	588	0.028	0	0	0	...			
<i>Andira inermis</i>	T	n	n	n	318	307	306	-0.004	42	36	33	-0.027			
<i>Annona acuminata</i>	S	n	W	n	509	526	568	0.012	0	0	0	...			
<i>Annona hayesii</i>	U	...	n	n	1	1	1	0.000	0	0	0	...			
<i>Annona spraguei</i>	M	C	n	n	57	71	143	0.108	18	18	22	0.023			
<i>Apeiba hybrid</i>	T	1	1	0	...	1	1	0	...			
<i>Apeiba membranacea</i>	T	...	n	n	389	342	330	-0.019	238	226	238	0.000			
<i>Apeiba tibourbou</i>	M	...	n	n	46	39	33	-0.039	26	23	22	-0.019			
<i>Aphelandra sinclairiana</i>	S	...	W	S	13	10	7	-0.077	0	0	0	...			
<i>Aphonia seibertii</i>	S	1	1	4	0.146	0	0	0	...			
<i>Ardisia barlettii</i>	U	...	n	n	3	3	1	-0.115	0	0	0	...			
<i>Ardisia fendleri</i>	S	...	n	S	77	80	85	0.011	1	1	0	...			
<i>Ardisia guianensis</i>	S	...	W	n	21	15	17	-0.026	0	0	0	...			
<i>Aspidosperma cruentata</i>	T	n	n	S	451	471	485	0.008	48	53	55	0.015			
<i>Astrocaryum standleyanum</i>	M	...	W	n	248	232	222	-0.013	233	225	217	-0.008			
<i>Astronium graveolens</i>	U	...	n	n	65	59	66	0.001	35	35	35	0.000			
<i>Bactris barronis</i>	U	...	n	n	112	57	23	-0.184	0	0	0	...			
<i>Bactris colonata</i>	S	...	n	S	240	83	23	-0.267	0	0			
<i>Bactris coloradensis</i>	U	...	n	n	40	18	6	-0.222	0	0	0	...			

<i>Bactris major</i>	U	n	W	n	476	362	224	-0.090	0	0	0	...
<i>Banara guianensis</i>	U	...	W	S	0	0	4	...	0	0	0	...
<i>Belischmedia pendula</i>	T	n	n	n	2376	2671	2750	0.017	308	303	295	-0.004
<i>Bertiera guianensis</i>	S	...	n	n	2	2	1	-0.072	0	0	0	...
<i>Brosimum alicastrum</i>	T	n	n	n	865	897	925	0.007	183	179	183	0.000
<i>Brosimum guianense</i>	T	...	n	n	1	1	1	0.000	1	1	1	0.000
<i>Calophyllum longifolium</i>	T	n	n	S	649	722	893	0.037	55	54	60	0.010
<i>Capparis frondosa</i>	S	n	n	n	3536	3676	3550	0.000	0	2	1	...
<i>Casarea aculeata</i>	U	n	n	n	467	476	480	0.003	24	26	23	-0.004
<i>Casarea arborea</i>	T	C	n	n	21	23	21	0.000	2	3	3	0.052
<i>Casarea guianensis</i>	U	...	n	n	12	14	16	0.032	0	0	0	...
<i>Casarea sylvestris</i>	M	n	n	n	248	232	211	-0.019	72	67	58	-0.024
<i>Cassipourea elliptica</i>	M	n	W	n	765	850	945	0.025	67	72	78	0.018
<i>Cavanillesia platanifolia</i>	T	...	n	S	22	21	22	0.000	21	21	21	0.000
<i>Cecropia insignis</i>	T	C	n	n	517	447	405	-0.028	280	249	261	-0.008
<i>Cecropia obtusifolia</i>	M	C	W	n	61	38	48	-0.028	38	24	22	-0.064
<i>Cedrela odorata</i>	T	...	n	S	5	2	9	0.068	2	2	2	0.000
<i>Ceiba pentandra</i>	T	...	n	n	72	67	61	-0.019	42	40	37	-0.014
<i>Celtis schippii</i>	M	...	n	S	182	166	148	-0.023	42	38	32	-0.031
<i>Cespedeza macrophylla</i>	T	n	7	5	5	-0.037	3	2	2	-0.046
<i>Cestrum megalophyllum</i>	S	...	W	S	309	237	157	-0.077	0	0	0	...
<i>Chamaedorea tepejilote</i>	S	...	n	S	32	22	16	-0.080	0	0	0	...
<i>Chamguava schippii</i>	U	n	n	n	195	240	289	0.048	1	1	2	0.085
<i>Chamarrhis parviflora</i>	T	...	n	n	2	2	1	-0.086	0	0	1	...
<i>Chrysochlamys eclipes</i>	S	n	n	S	460	438	423	-0.009	2	5	3	0.043
<i>Chrysophyllum argenteum</i>	T	C	n	n	423	477	683	0.056	75	84	84	0.013
<i>Chrysophyllum cainito</i>	T	C	W	n	70	80	109	0.051	21	23	25	0.019
<i>Citidemia dentata</i>	S	...	W	n	8	14	17	0.094	0	0	0	...
<i>Citidemia octona</i>	S	...	n	n	3	6	9	0.124	0	0	0	...
<i>Citidemia septuflneria</i>	S	1	2	0	...	0	0	0	...
<i>Cnemidaria petiolata</i>	S	8	3	1	-0.252	0	0	0	...
<i>Coccoloba coronata</i>	M	n	n	n	181	191	185	0.002	22	21	20	-0.011
<i>Coccoloba manzanillensis</i>	U	n	n	n	445	448	473	0.007	8	9	12	0.047
<i>Colubrina glandulosa</i>	T	...	n	n	2	1	1	-0.074	2	1	1	-0.074
<i>Conostegia cinamomea</i>	S	...	n	n	4	4	3	-0.030	0	0	0	...
<i>Cordia alliodora</i>	S	n	W	S	391	281	209	-0.071	0	0	0	...
<i>Cordia bicolor</i>	T	C	n	n	111	108	119	0.008	61	60	60	-0.001
<i>Cordia lasiocalyx</i>	M	C	n	n	715	758	1042	0.046	255	258	283	0.012
<i>Coussarea curvigenmia</i>	M	n	n	S	1698	1665	1688	0.000	420	442	441	0.005
<i>Coultaria hexandra</i>	U	n	n	n	1502	1663	1963	0.032	40	46	57	0.042
<i>Croton bilbergianus</i>	U	1	1	1	0.000	0	0	0	...
<i>Croton bilbergianus</i>	U	C	W	n	620	619	1012	0.059	67	67	75	0.013
<i>Cupania cinerea</i>	M	...	n	n	6	7	12	0.083	1	1	1	0.000

Table 1. (cont.)

Species	Codes			Stems ≥ 10 mm dbh			Stems ≥ 100 mm dbh			Rate 1982-1990	
	G	C	S	1982	1985	1990	1982	1985	1990		
											Family
<i>Cupania latifolia</i>	T	...	n	S	54	48	51	8	9	10	0.025
<i>Cupania refescens</i>	T	n	W	n	55	71	96	5	6	6	0.022
<i>Cupania sylvatica</i>	U	n	n	n	962	1040	1146	28	30	41	0.045
<i>Cyphomandra hartwegii</i>	S	...	n	n	5	4	3	0	0	0	...
<i>Dendropanax arboreus</i>	T	...	n	n	18	18	16	11	11	11	0.000
<i>Dendropanax stenodontus</i>	T	...	n	n	159	144	121	96	93	84	-0.016
<i>Desmopsis panamensis</i>	U	n	n	n	11735	12141	12167	17	15	12	-0.040
<i>Diospyros aranthifolia</i>	M	n	n	n	48	54	70	9	12	14	0.052
<i>Dipteryx panamensis</i>	T	...	n	n	56	52	53	33	33	34	0.003
<i>Drypetes standleyi</i>	T	n	n	S	2173	2266	2279	196	227	262	0.031
<i>Elaeis oleifera</i>	M	...	W	...	22	21	21	22	21	21	-0.005
<i>Enterobium schomburgkii</i>	T	...	n	n	13	12	13	2	2	2	0.000
<i>Eyleria costaricensis</i>	U	...	n	S	289	236	185	47	46	30	-0.051
<i>Erythroxylum macrophyllum</i>	M	n	n	n	327	308	300	18	20	16	-0.013
<i>Erythroxylum panamense</i>	U	n	W	n	104	105	111	0	0	0	...
<i>Eugenia coloradensis</i>	T	n	n	n	803	840	832	78	79	83	0.007
<i>Eugenia galalonensis</i>	U	n	n	n	961	1157	1379	21	25	16	-0.031
<i>Eugenia nesiotica</i>	M	n	n	n	509	529	539	48	48	53	0.011
<i>Eugenia oerstediana</i>	M	n	n	n	2088	2208	2346	133	140	159	0.020
<i>Faramaea occidentalis</i>	U	n	n	n	23460	25154	26912	1228	1402	1652	0.034
<i>Ficus bullenii</i>	1	1	1	1	1	1	0.000
<i>Ficus citrifolia</i>	H	...	n	n	0	0	1	0	0	0	...
<i>Ficus colubrinæ</i>	n	n	1	1	1	1	1	1	0.000
<i>Ficus costaricana</i>	H	...	W	n	16	11	7	8	7	7	-0.014
<i>Ficus insipida</i>	T	...	W	n	9	8	7	3	3	3	0.000
<i>Ficus maxima</i>	T	...	W	S	11	11	10	5	6	3	-0.059
<i>Ficus obtusifolia</i>	H	...	W	n	12	10	8	10	8	8	-0.023
<i>Ficus pertusa</i>	H	...	n	n	0	0	1	0	0	0	...
<i>Ficus popenoei</i>	H	...	W	n	9	8	7	7	6	5	-0.039
<i>Ficus tonduzii</i>	M	...	n	S	66	61	45	42	37	30	-0.039
<i>Ficus trigonata</i>	H	...	n	S	8	5	5	6	4	5	-0.019
<i>Ficus yuponensis</i>	T	...	n	n	6	6	6	6	6	6	0.000
<i>Garcinia intermedia</i>	M	n	n	n	3654	4023	4300	77	75	81	0.005
<i>Garcinia madruno</i>	M	n	n	n	652	682	524	23	26	24	0.005
<i>Genipa americana</i>	T	...	W	n	91	88	84	20	20	21	0.005
<i>Geonoma interrupta</i>	U	19	14	3	0	0	0	...
<i>Guaphira standleyanum</i>	T	...	n	n	230	208	191	90	91	99	0.010

<i>Guarea grandifolia</i>	T	...	n	n	55	57	62	0.014	10	10	10	0.000
<i>Guarea guidonia</i>	M	n	W	S	1781	1829	1968	0.011	370	363	376	0.001
<i>Guarea sp. nov.</i>	M	n	n	n	1558	1492	1378	-0.014	105	95	83	-0.026
<i>Guatteria dumetorum</i>	T	n	n	S	1587	1529	1472	-0.008	285	280	277	-0.003
<i>Guazuma ulmifolia</i>	T	...	W	n	55	50	52	-0.006	28	30	34	0.021
<i>Guettarda foliacea</i>	U	n	n	n	382	382	369	-0.004	78	83	80	0.002
<i>Gustavia superba</i>	M	C	n	n	881	821	828	-0.007	637	642	650	0.002
<i>Hamelia axillaris</i>	S	n	W	n	114	92	107	-0.007	1	0	0	...
<i>Hamelia patens</i>	n	n	0	2	2	...	0	0	0	...
<i>Hamepa appendiculata</i>	M	...	n	n	76	49	40	-0.076	41	33	25	-0.058
<i>Hasseltia floribunda</i>	M	n	W	n	1147	1019	866	-0.033	262	256	249	-0.006
<i>Heisteria acuminata</i>	U	n	n	n	105	107	108	0.003	5	6	7	0.039
<i>Heisteria concinna</i>	M	n	n	n	888	964	988	0.012	246	256	275	0.012
<i>Herrania purpurea</i>	U	C	n	n	522	531	548	0.005	0	0	0	...
<i>Hirtella americana</i>	T	...	n	n	50	47	47	-0.006	8	8	6	-0.031
<i>Hirtella triandra</i>	M	n	n	S	4144	4661	5025	0.022	516	554	616	0.020
<i>Hura crepitans</i>	T	...	W	n	127	119	117	-0.009	100	97	100	0.000
<i>Hybanthus prunifolius</i>	S	n	n	n	39872	41154	40511	0.001	3	3	1	-0.125
<i>Hyevontima alcheornoides</i>	T	C	W	n	101	100	99	-0.002	44	42	42	-0.005
<i>Inga acuminata</i>	U	n	n	n	07	250	300	0.042	11	14	20	0.068
<i>Inga cocleensis</i>	M	C	n	n	219	219	232	0.007	39	39	46	0.020
<i>Inga jagifolia</i>	T	...	W	n	58	60	70	0.022	8	7	6	-0.031
<i>Inga goldmanii</i>	T	n	n	n	498	468	462	-0.008	62	51	51	-0.022
<i>Inga marginata</i>	T	n	n	S	913	810	741	-0.023	81	74	91	0.013
<i>Inga minutula</i>	T	...	n	S	12	12	11	-0.010	4	3	2	-0.081
<i>Inga mucuna</i>	T	...	n	n	1	1	1	0.000	0	0	0	...
<i>Inga pezi-zifera</i>	T	n	n	S	208	202	184	-0.013	25	15	13	-0.068
<i>Inga punctata</i>	T	...	W	S	17	18	20	0.018	8	7	7	-0.015
<i>Inga quaternata</i>	M	n	n	S	735	742	731	0.000	34	40	48	0.038
<i>Inga ruiziana</i>	T	...	n	n	80	70	62	-0.030	10	6	10	0.000
<i>Inga sapindoides</i>	M	C	n	n	391	354	327	-0.021	67	67	75	0.013
<i>Inga umbellifera</i>	T	...	n	n	22	24	25	0.015	12	12	14	0.018
<i>Jacaranda copaia</i>	M	n	n	n	942	997	1040	0.011	21	17	12	-0.065
<i>Koanophyllon wetmorei</i>	T	C	n	n	342	319	298	-0.016	224	230	240	0.008
<i>Lacistema aggregatum</i>	S	...	n	S	15	12	12	-0.029	0	0	0	...
<i>Lacistema panamensis</i>	U	n	n	n	1557	1661	1694	0.010	43	40	37	-0.017
<i>Laetia procer</i>	M	...	W	n	91	91	98	0.008	36	37	42	0.018
<i>Laetia thamnia</i>	T	...	n	n	24	23	30	0.027	9	9	11	0.024
<i>Lafoesia puniceifolia</i>	U	n	n	n	514	519	542	0.006	15	17	25	0.061
<i>Leandra dichotoma</i>	T	...	W	n	8	7	6	-0.035	4	4	5	0.027
<i>Licania hypoleuca</i>	S	...	n	n	0	1	1	...	0	0	0	...
<i>Licania platypus</i>	M	n	n	n	120	121	133	0.012	19	16	15	-0.027
	T	n	n	S	253	293	315	0.024	9	9	10	0.011

Table 1. (cont.)

Species	Codes						Stems ≥ 10 mm dbh			Stems ≥ 100 mm dbh			Rate 1982-1990	
	G	C	W	S	Family	Rate 1982-1990	1982	1985	1990	1982	1985	1990		Rate 1982-1990
<i>Lindackeria laurina</i>	M	...	W	n	Flacourtiaceae	-0.023	109	97	89	85	78	69	-0.023	
<i>Lonicocarpus latifolia</i>	T	C	n	n	Leguminosae	0.006	842	846	889	147	137	124	-0.020	
<i>Lopimia dasypetala</i>	S	Malvaceae	...	1	1	0	0	0	0	...	
<i>Lozania pittieri</i>	U	Lacisternaceae	-0.117	18	15	7	3	2	0	...	
<i>Luehea seemannii</i>	T	C	W	n	Tiliaceae	0.020	188	190	225	87	93	94	0.008	
<i>Lycianthes maxonii</i>	S	Solanaceae	...	1	1	0	0	0	0	...	
<i>Maclura tinctoria</i>	T	Moraceae	-0.086	2	2	1	0	1	1	...	
<i>Macrocnemum glabrescens</i>	M	C	n	S	Rubiaceae	0.010	96	101	106	24	25	25	0.004	
<i>Malmea sp. nov.</i>	M	n	n	n	Annonaceae	0.035	276	320	372	14	15	17	0.023	
<i>Malpighia romeroana</i>	S	n	n	n	Malpighiaceae	0.012	54	64	60	0	0	0	...	
<i>Maquira costaricana</i>	M	n	n	n	Moraceae	0.006	1418	1445	1503	223	200	176	-0.027	
<i>Margaritaria nobilis</i>	U	...	n	n	Euphorbiaceae	0.144	1	2	4	1	1	1	0.000	
<i>Marila laxiflora</i>	M	...	n	S	Guttiferae	0.000	22	26	22	10	11	11	0.010	
<i>Mytenus schippii</i>	M	...	n	n	Celastraceae	0.008	82	82	88	20	19	20	0.000	
<i>Miconia affinis</i>	U	C	n	n	Melastomataceae	0.020	371	394	439	4	2	9	0.098	
<i>Miconia argentea</i>	M	C	W	n	Melastomataceae	0.062	531	678	902	45	50	56	0.025	
<i>Miconia elata</i>	U	...	n	n	Melastomataceae	-0.015	34	31	30	2	3	2	0.000	
<i>Miconia hondurensis</i>	U	...	W	n	Melastomataceae	0.025	30	29	37	7	4	6	-0.018	
<i>Miconia impatiolaris</i>	U	...	W	S	Melastomataceae	0.031	6	5	8	0	0	0	...	
<i>Miconia nervosa</i>	S	C	n	n	Melastomataceae	-0.014	359	294	320	0	0	0	...	
<i>Miconia prasina</i>	U	...	n	n	Melastomataceae	...	0	0	2	0	0	0	...	
<i>Mouriri myrtilloides</i>	S	n	n	n	Melastomataceae	0.011	6948	7712	7618	0	1	0	...	
<i>Myrcia gatunensis</i>	U	n	n	n	Myrtaceae	0.040	40	49	56	2	2	4	0.082	
<i>Myrspermum frutescens</i>	T	...	n	n	Leguminosae	-0.021	28	24	23	9	9	10	0.011	
<i>Nectandra cissiflora</i>	T	n	n	n	Lauraceae	-0.013	349	339	314	23	20	27	0.019	
<i>Nectandra globosa</i>	M	C	W	S	Lauraceae	0.001	117	113	119	12	10	13	0.009	
<i>Nectandra purpurea</i>	M	n	n	n	Lauraceae	0.001	80	77	81	4	2	3	-0.035	
<i>Nectandra sp. nov. 1</i>	T	...	n	S	Lauraceae	-0.044	9	9	6	1	0	0	...	
<i>Nectandra sp. nov. 3</i>	Lauraceae	...	1	1	0	0	0	0	...	
<i>Neea amplifolia</i>	S	n	n	n	Nyctaginaceae	0.011	62	71	68	0	0	0	...	
<i>Ochroma pyramidale</i>	M	...	n	n	Bombacaceae	0.040	5	9	7	3	5	6	0.082	
<i>Ocotea cernua</i>	M	...	n	n	Lauraceae	-0.004	346	337	332	24	28	26	0.009	
<i>Ocotea oblonga</i>	T	n	n	n	Lauraceae	-0.014	215	184	190	33	31	27	-0.022	
<i>Ocotea puberula</i>	T	n	n	n	Lauraceae	-0.023	269	243	222	9	10	16	0.069	
<i>Ocotea whitlei</i>	T	n	n	S	Lauraceae	-0.042	1130	941	769	167	170	176	0.005	
<i>Onocarpus mapourea</i>	M	...	n	n	Palmae	0.000	1790	1712	1802	752	746	754	0.000	
<i>Olmedia aspera</i>	U	n	n	S	Moraceae	-0.051	442	376	279	50	42	33	-0.046	

<i>Ormosia amazonica</i>	T	1	2	2	0.078	1	1	1	0.000
<i>Ormosia croatii</i>	T	n	n	S	52	56	71	0.034	4	3	4	0.000
<i>Ormosia macrocalyx</i>	T	...	n	n	4	4	4	0.000	4	4	4	0.000
<i>Ouratea lucens</i>	S	n	n	n	1122	1240	1266	0.014	0	0	0	...
<i>Pallicourea guianensis</i>	S	C	W	n	377	663	1475	0.165	0	0	0	...
<i>Pentagonia macrophylla</i>	U	n	n	S	568	513	453	-0.026	2	2	2	0.000
<i>Pereba xanthochyma</i>	M	n	n	n	268	268	273	0.002	13	13	19	0.047
<i>Phoebe cinnamomifolia</i>	T	C	n	n	88	83	87	-0.001	18	18	19	0.006
<i>Picramnia latifolia</i>	U	n	n	S	1169	1177	1174	0.000	38	39	42	0.011
<i>Piper aequale</i>	S	n	n	S	219	158	83	-0.113	0	0	0	...
<i>Piper arboreum</i>	U	...	n	S	107	83	60	-0.069	0	0	0	...
<i>Piper carrilloanum</i>	S	12	10	4	-0.121	0	0	0	...
<i>Piper cordulatum</i>	S	n	n	n	3149	3713	1777	-0.069	1	0	0	...
<i>Piper cuberatum</i>	S	...	W	S	120	65	53	-0.099	0	0	0	...
<i>Piper imperiale</i>	U	9	3	3	-0.126	0	0	0	...
<i>Piper perlasense</i>	S	n	n	S	110	117	68	-0.052	0	0	0	...
<i>Piper reticulatum</i>	U	n	W	S	180	169	149	-0.022	9	9	9	0.000
<i>Pithecellobium macradenium</i>	T	...	n	n	1	1	1	0.000	1	1	1	0.000
<i>Pithecellobium rufescens</i>	M	...	n	n	2	2	2	0.000	0	0	0	...
<i>Platymiscium pinnatum</i>	T	n	n	n	256	248	266	0.004	71	69	64	-0.012
<i>Platypodium elegans</i>	R	C	n	n	170	157	159	-0.007	58	49	45	-0.029
<i>Pochota quinta</i>	T	...	n	n	2	1	1	-0.083	2	1	1	-0.083
<i>Pochota sessilis</i>	T	25	24	23	-0.009	9	11	11	0.022
<i>Posoqueria latifolia</i>	M	...	n	n	77	80	77	0.000	14	15	14	0.000
<i>Poulsenia armata</i>	T	n	n	S	3430	2686	2126	-0.054	923	857	801	-0.016
<i>Pourouma bicolor</i>	T	...	n	S	40	43	43	0.009	11	12	15	0.039
<i>Pouteria fossicola</i>	T	...	n	S	3	3	3	0.000	2	2	2	0.000
<i>Pouteria reticulata</i>	T	n	n	n	1652	1718	1766	0.007	157	170	192	0.023
<i>Pouteria stipitata</i>	M	...	n	n	61	61	62	0.001	33	31	29	-0.015
<i>Prioria copajfera</i>	T	n	n	n	1356	1408	1442	0.007	279	309	336	0.022
<i>Protium costaricense</i>	M	n	n	S	913	860	870	-0.005	110	101	105	-0.005
<i>Protium panamense</i>	M	n	n	n	2715	2844	3029	0.013	65	51	57	-0.015
<i>Protium sp. nov.</i>	M	...	n	S	0	1	3	...	0	0	0	...
<i>Protium tenuifolium</i>	M	n	n	n	2664	2905	3081	0.016	354	353	362	0.002
<i>Pseudobombax septenatum</i>	T	...	W	n	22	15	18	-0.022	11	9	8	-0.035
<i>Psidium friedrichsthalianum</i>	U	...	W	S	39	42	47	0.020	4	4	4	0.000
<i>Psychotria acuminata</i>	S	...	W	n	8	11	16	0.086	0	0	0	...
<i>Psychotria brachiata</i>	S	0	0	1	...	0	0	0	...
<i>Psychotria chagrensis</i>	S	n	n	S	13	17	15	0.017	0	0	0	...
<i>Psychotria deflexa</i>	S	n	n	n	88	77	60	-0.047	0	0	0	...
<i>Psychotria furcata</i>	S	...	W	n	5	1	2	0.000	0	0	0	...
<i>Psychotria graciliflora</i>	S	...	W	n	10	14	44	0.185	0	0	0	...
<i>Psychotria granadensis</i>	S	...	n	n	7	5	4	-0.065	0	0	0	...

Table 1. (cont.)

Species	Codes						Stems ≥ 10 mm dbh				Stems ≥ 100 mm dbh				Rate	
	G	C	W	S	Family	Rate	1982	1985	1990	Rate	1982	1985	1990	Rate	1982-1990	
																1982-1990
<i>Psychotria grandis</i>	U	n	W	S	Rubiaceae	104	97	81	-0.027	2	2	2	0.000			
<i>Psychotria horizontalis</i>	S	n	n	n	Rubiaceae	6168	6446	5920	-0.004	0	0	0	...			
<i>Psychotria limonensis</i>	S	...	n	n	Rubiaceae	15	17	19	0.027	0	0	0	...			
<i>Psychotria marginata</i>	S	n	n	n	Rubiaceae	582	691	696	0.021	0	0	0	...			
<i>Psychotria pittieri</i>	S	...	n	S	Rubiaceae	4	3	3	-0.031	0	0	0	...			
<i>Psychotria racemosa</i>	S	...	n	n	Rubiaceae	1	2	2	0.086	0	0	0	...			
<i>Pterocarpus helizensis</i>	T	...	n	n	Leguminosae	9	8	3	-0.125	1	1	0	...			
<i>Pterocarpus rohrii</i>	T	n	n	n	Leguminosae	1577	1629	1705	0.009	136	103	88	-0.056			
<i>Quararibea asterolepis</i>	T	n	n	n	Bombacaceae	2394	2379	2348	-0.002	703	694	699	0.000			
<i>Quassia amara</i>	U	...	n	n	Simaroubaceae	155	151	147	-0.005	7	8	8	0.014			
<i>Randia armata</i>	U	n	n	n	Rubiaceae	1128	1155	1143	0.001	228	234	241	0.006			
<i>Randia formosa</i>	U	...	n	n	Rubiaceae	4	4	4	0.000	0	0	0	...			
<i>Rinorea sylvatica</i>	S	n	n	n	Violaceae	2571	2615	2596	0.001	1	0	0	...			
<i>Sapium acutiparum</i>	T	...	W	n	Euphorbiaceae	47	41	40	-0.018	23	20	16	-0.040			
<i>Sapium</i> sp. nov.	T	Euphorbiaceae	3	3	3	0.000	3	3	3	0.000			
<i>Scheelea zonenis</i>	M	...	W	n	Palmae	44	40	36	-0.023	44	40	36	-0.023			
<i>Schefflera morototoni</i>	T	Araliaceae	1	1	0	...	1	1	0	...			
<i>Schizolobium paralylybium</i>	T	...	n	n	Leguminosae	13	13	16	0.026	1	1	1	0.000			
<i>Senna darimensis</i>	S	C	n	n	Leguminosae	205	138	116	-0.065	1	2	2	0.080			
<i>Simarouba amara</i>	T	n	n	n	Simaroubaceae	1240	1250	1292	0.004	247	255	269	0.010			
<i>Siparuna guianensis</i>	M	...	n	S	Monimiaceae	37	34	32	-0.018	16	15	15	-0.008			
<i>Siparuna pauciflora</i>	U	n	n	S	Monimiaceae	431	354	316	-0.035	24	26	18	-0.033			
<i>Sloanea terniflora</i>	T	n	n	n	Elaeocarpaceae	601	591	575	-0.005	85	85	80	-0.006			
<i>Socratea exorrhiza</i>	M	n	n	n	Palmae	812	737	684	-0.021	374	357	336	-0.013			
<i>Solanum arboreum</i>	S	Solanaceae	1	1	0	...	0	0	0	...			
<i>Solanum asperum</i>	S	...	W	n	Solanaceae	0	0	4	...	0	0	0	...			
<i>Solanum hayesii</i>	M	C	n	S	Solanaceae	125	89	77	-0.058	40	25	13	-0.136			
<i>Solanum stephmarkii</i>	S	...	n	n	Solanaceae	7	6	2	-0.145	0	0	0	...			
<i>Sorocea affinis</i>	S	n	n	n	Moraceae	3302	3371	3302	0.000	47	44	30	-0.051			
<i>Spaethea membranacea</i>	U	...	n	n	Malpighiaceae	19	17	15	-0.028	8	8	7	-0.015			
<i>Spondias mombin</i>	T	C	W	n	Anacardiaceae	63	67	101	0.057	24	23	30	0.026			
<i>Spondias radlkoferi</i>	T	C	n	n	Anacardiaceae	192	164	202	0.006	55	56	57	0.004			
<i>Stemmadenia grandiflora</i>	U	Apocynaceae	1	1	0	...	1	1	0	...			
<i>Sterculia apetala</i>	T	...	W	n	Sterculiaceae	68	53	65	-0.005	25	24	27	0.008			
<i>Stylopne standleyi</i>	S	n	W	n	Myrsinaceae	713	735	751	0.006	1	1	0	...			
<i>Swartzia simplex</i> (var. <i>grandiflora</i>)	U	n	n	n	Leguminosae	2255	2422	2574	0.015	198	203	212	0.007			

<i>Swartzia simplex</i> (var. <i>ochracea</i>)	U	n	n	n	2701	2821	2857	0.006	104	112	123	0.019
<i>Symphonia globulifera</i>	T	n	n	S	188	178	175	-0.008	46	38	30	-0.048
<i>Tabebuia guayacan</i>	T	...	n	n	76	74	73	-0.004	30	28	28	-0.007
<i>Tabebuia rosea</i>	T	n	W	n	316	300	318	0.000	81	75	71	-0.014
<i>Tabernaemontana arborea</i>	T	n	W	n	1287	1328	1421	0.011	293	302	303	0.004
<i>Tachigali versicolor</i>	T	n	n	n	2923	2980	3095	0.006	86	82	88	0.002
<i>Talisia nervosa</i>	U	n	n	n	813	819	802	-0.001	0	0	0	...
<i>Talisia princeps</i>	M	n	n	n	623	638	673	0.008	7	9	5	-0.038
<i>Terminalia amazonica</i>	T	...	W	n	62	60	59	-0.005	28	28	28	0.000
<i>Terminalia oblonga</i>	T	...	n	n	92	90	98	0.007	43	42	46	0.007
<i>Temstroemia tepezapote</i>	U	...	n	n	1	1	1	0.000	0	0	1	...
<i>Tetragastris panamensis</i>	T	n	n	n	3233	3699	4085	0.026	318	323	353	0.012
<i>Tetralylacium johansensii</i>	T	...	W	n	10	7	7	-0.039	8	7	6	-0.032
<i>Theobroma cacao</i>	U	...	n	S	22	20	21	-0.005	12	14	13	0.009
<i>Thevetia abouai</i>	U	n	W	S	107	97	97	-0.011	2	1	1	-0.078
<i>Tocoyena pittieri</i>	M	...	W	S	6	8	7	0.017	4	5	5	0.026
<i>Trattinnickinia aspera</i>	T	...	n	n	112	96	82	-0.036	50	47	48	-0.004
<i>Trema micrantha</i>	M	...	n	n	32	23	21	-0.051	21	18	15	-0.041
<i>Trichanthera gigantea</i>	U	...	n	n	12	9	8	-0.048	3	3	2	-0.047
<i>Trichilia pallida</i>	M	n	W	n	567	574	591	0.004	76	75	80	0.006
<i>Trichilia tuberculata</i>	T	n	n	n	12927	13166	13298	0.003	2022	1902	1783	-0.014
<i>Trichospermum galeottii</i>	M	...	n	n	7	4	2	-0.154	6	4	2	-0.135
<i>Triplaris camigiana</i>	M	n	W	n	371	342	319	-0.016	125	127	145	0.016
<i>Trophis racemosa</i>	M	n	n	S	312	323	325	0.004	54	48	38	-0.040
<i>Turpinia occidentalis</i>	T	...	n	n	153	113	85	-0.069	69	56	57	-0.022
<i>Unonopsis pittieri</i>	M	n	n	S	771	785	787	0.002	136	148	160	0.019
<i>Urea baccifera</i>	S	...	n	S	2	5	17	0.240	0	0	0	...
<i>Viola sebifera</i>	M	n	n	n	2406	2276	2086	-0.016	607	588	607	0.000
<i>Viola sp. nov.</i>	T	...	n	S	58	53	53	-0.010	21	20	23	0.010
<i>Viola surinamensis</i>	T	...	n	S	300	259	239	-0.025	174	163	158	-0.010
<i>Vismia baccifera</i>	U	n	W	n	75	77	80	0.007	1	3	2	0.075
<i>Vismia bilbergiana</i>	U	3	4	3	0.000	0	0	0	...
<i>Vismia macrophylla</i>	M	1	0	0	...	1	0	0	...
<i>Vochysia ferruginea</i>	T	...	W	n	35	29	29	-0.022	10	11	11	0.011
<i>Xylofia macrantha</i>	M	n	n	S	817	916	1044	0.027	79	99	128	0.054
<i>Xylosma chloranthum</i>	U	1	1	0	...	0	0	0	...
<i>Xylosma obigandrum</i>	S	n	n	n	183	169	149	-0.024	1	1	1	0.000
<i>Zanthosyllum belizense</i>	T	C	n	n	220	252	237	0.008	103	108	140	0.036
<i>Zanthosyllum panamense</i>	T	C	n	n	297	238	227	-0.031	83	70	66	-0.027
<i>Zanthosyllum procerum</i>	M	C	n	n	207	212	170	-0.023	26	28	35	0.034
<i>Zanthosyllum setulosum</i>	M	1	1	1	0.000	1	1	1	0.000
<i>Zuelania guianensis</i>	M	...	W	n	40	41	40	0.000	13	14	11	-0.018

Table 2. Species whose population of stems ≥ 10 mm changed at a rate $\geq 5\%$ per year. The four columns of species characteristics are the same as those given in Table 1 (growth form, colonizing, swamp and slope status). Species are ordered from those with the fastest shrinking populations to the fastest growing; the line separates growing from shrinking populations.

Species	G	C	W	S	Population		Rate of population change		
					1982	1990	1982–1985	1985–1990	1982–1990
<i>Piper aequale</i>	S	n	n	S	219	83	-0.0984	-0.1223	-0.1131
<i>Piper culebratum</i>	S	...	W	S	120	53	-0.2036	-0.0391	-0.0992
<i>Chamaedorea tepejilote</i>	S	...	n	S	32	16	-0.1128	-0.0600	-0.0803
<i>Cestrum megalophyllum</i>	S	...	W	S	309	157	-0.0769	-0.0783	-0.0777
<i>Hampea appendiculata</i>	M	...	n	n	76	40	-0.1363	-0.0388	-0.0760
<i>Acalypha diversifolia</i>	S	C	W	n	1568	827	-0.0742	-0.0717	-0.0727
<i>Acalypha macrostachya</i>	U	C	n	n	80	45	-0.0626	-0.0762	-0.0714
<i>Conostegia bracteata</i>	S	n	W	S	391	209	-0.0931	-0.0563	-0.0711
<i>Piper cordulatum</i>	S	n	n	n	3149	1777	0.0545	-0.1407	-0.0693
<i>Turpinia occidentalis</i>	T	...	n	n	153	85	-0.0937	-0.0539	-0.0690
<i>Piper arboreum</i>	U	...	n	S	107	60	-0.0798	-0.0624	-0.0690
<i>Senna dariensis</i>	S	S	n	n	205	116	-0.1171	-0.0328	-0.0657
<i>Solanum hayesii</i>	M	C	n	S	125	77	-0.1121	-0.0273	-0.0582
<i>Poulsenia armata</i>	T	n	n	S	3430	2126	-0.0703	-0.0441	-0.0545
<i>Piper perlasense</i>	S	n	n	S	110	68	0.0162	-0.1029	-0.0529
<i>Erythrina costaricana</i>	U	...	n	S	289	185	-0.0622	-0.0464	-0.0525
<i>Trema micrantha</i>	M	...	n	n	32	21	-0.1167	-0.0172	-0.0519
<i>Olmedia aspera</i>	U	n	n	S	442	279	-0.0434	-0.0564	-0.0510
<i>Chrysophyllum cainito</i>	T	C	W	n	70	109	0.0399	0.0579	0.0510
<i>Chrysophyllum argenteum</i>	T	C	n	n	423	683	0.0366	0.0681	0.0560
<i>Spondias mombin</i>	T	C	W	n	63	101	0.0207	0.0785	0.0575
<i>Croton billbergianus</i>	U	C	W	n	620	1012	-0.0005	0.0944	0.0590
<i>Miconia argentea</i>	M	C	W	n	531	902	0.0764	0.0542	0.0626
<i>Cupania refescens</i>	T	n	W	n	55	96	0.0775	0.0578	0.0654
<i>Annona spraguei</i>	M	C	n	n	55	143	0.0680	0.1328	0.1082
<i>Palicourea guianensis</i>	S	C	w	n	377	1475	0.1861	0.1533	0.1654
<i>Psychotria graciliflora</i>	S	...	W	n	10	44	0.1231	0.2176	0.1853

(38%) changed by $< 1\%$ per year, and 11 (8%) changed by more than 5% per year (eight of the latter were declines and three increases). The fastest rate of change in the larger size class was *Inga acuminata*, which increased from 11 to 20 stems; a more common species increasing nearly as rapidly was *Xylopia macrantha*, whose population rose from 79 to 128 stems. The greatest decline was in *Solanum hayesii*, which had 40 stems in 1982 but just 13 in 1990. A more abundant species, *Pterocarpus rohrii*, fell from 136 to 83 stems. The mean rate of population change among 136 species in the large size class was -0.32% , and the mean rate of absolute change was 1.93% . These are not significantly different from rates for stems ≥ 10 mm (t-test).

Within-species consistency in population change

Rates of change during 1982–1985 and 1985–1990 were consistent within species (Figure 2A; the correlation is highly significant: $r^2 = 0.266$, $P < 0.0001$ for stems ≥ 10 mm dbh; $r^2 = 0.246$, $P < 0.0001$ for stems ≥ 100 mm). There were




Figure 2. Scatter plots of population change, with one point plotted for each species. The dotted lines are diagonal regressions, shown only to indicate the direction of trends. (A) Population change for stems ≥ 10 mm dbh, plotting 1985–1990 rate *vs* 1982–1985; 219 species included. (B) Population change over 1982–1990, plotting rate for stems ≥ 100 mm dbh *vs* rate for stems ≥ 10 mm dbh; 136 species included.

Table 3. Numbers of species with increasing and decreasing populations, by colonizing status, slope status and growth form. 'Slope' means slope-specialists that were not colonizers, 'colonizer' means colonizing species that were not slope-specialists and 'neither' means species that were neither slope-specialists nor colonizers (unlike Figure 1, this latter category does not include species for which information was missing). The four slope-colonizer species are not included. Asterisks between the increasing and decreasing columns indicate a significant difference in the fraction of species increasing between a given group and the 'neither group' (** indicating $P < 0.01$ and * $P < 0.05$). The final column gives the mean rate of population change for each species group.

Growth form	Status	Number of populations			Mean rate of change
		Increasing	Decreasing	No change	
Large trees	Slope	5	6	0	-0.004 ± 0.030
	Neither	15	7	0	0.005 ± 0.017
	Colonizer	8	8	0	0.003 ± 0.032
Mid-sized trees	Slope	5	3	0	0.008 ± 0.012
	Neither	17	11	0	0.001 ± 0.020
	Colonizer	4	4	0	0.018 ± 0.049
Understorey trees	Slope	1	**	6	-0.025 ± 0.016
	Neither	19	2	0	0.016 ± 0.016
	Colonizer	3	2	0	-0.002 ± 0.049
Shrubs	Slope	0	*	4	-0.062 ± 0.043
	Neither	10	5	1	-0.004 ± 0.024
	Colonizer	1	3	0	0.003 ± 0.011
All species	Slope	11	**	19	-0.013 ± 0.033
	Neither	61	25	1	0.005 ± 0.020
	Colonizer	16	*	17	0.006 ± 0.050

cases, though, where populations increased dramatically prior to 1985 then declined afterwards, or vice versa. For example, *Piper cordulatum* increased from 3149 to 3713 stems, then declined to 1777 (Table 1). Rate of change was also fairly consistent between the two size classes ($r^2 = 0.265$, $P < 0.0001$ for the 1982–1990 rate), but again there were exceptions (Figure 2B). For example, *Trichilia tuberculata*, the most abundant large tree in the plot, suffered a rather considerable decline among stems ≥ 100 mm dbh, but its population ≥ 10 mm dbh increased (Table 1).

Population change as a function of species characteristics

Slope status. Slope status was clearly associated with a species' probability of declining in abundance. Excluding all colonizing species (in order to separate the effect of that variable) there were 30 slope-specialists, 19 declining in abundance and 11 increasing. Of 87 non-slope species, 25 declined and 61 increased (Table 3). Thus, 29% of non-slope but 63% of slope-species declined in abundance ($\chi^2 = 11.1$, $df = 1$, $P < 0.01$). Sample sizes were augmented if colonizers were included, and the pattern remained: 63% of 52 slope species but 42% of 165 non-slope species declined in abundance.

The pattern did not hold for all growth forms, though, in fact the distinction between slope and non-slope species was due entirely to species of smaller stature – shrubs and treelets. Of this group, 91% of slope-species and only 19%

of non-slope species declined in abundance, with the difference statistically significant in both growth forms (Table 3). But in large and mid-sized trees, there was no such distinction, with 47% of slope and 36% of non-slope species declining in abundance and no significant difference in either growth form (Table 3).

Slope-specialists suffered the most impressive population declines. Of the 18 species declining more than 5% per year, 11 were slope-specialists (Figure 1A, Table 2). The four most rapid declines were slope-specialist shrubs (Table 2). Conversely, none of the nine species increasing more than 5% per year were slope-specialists (Table 2).

The poor performance of slope-specialists was not evident when considering stems above 100 mm dbh, because there were few treelets and shrubs included in this size class. For large and mid-sized trees, slope-specialists performed no worse than non-slope: six of 17 slope species decreased in abundance, whereas of non-slope species, 20 of 40 declined. Among treelets and shrubs, slope-specialists did suffer more declines than non-slope species (two out of three *vs* four out of 13) but the sample was far too small to evaluate statistically.

Colonizing species. Species designated as gap-colonizers performed somewhat worse than non-colonizing species, although not as poorly as slope species. Excluding slope-specialists (to isolate the effect of the colonization variable), 52% of 33 colonizing species and 29% of 87 non-colonizers declined in abundance ($\chi^2 = 5.3$, $P < 0.05$, Table 3). Each of the four growth forms showed a comparable pattern, with colonizers doing slightly worse than non-colonizers, but none was statistically significant by itself (Table 3).

Despite the fact that colonizers on average performed poorly, they were over-represented among rapidly increasing populations. Seven of the nine species increasing faster than 5% per year were colonizers, and only one was not (the other had an unknown colonization index). In contrast, of the 18 species decreasing faster than 5% per year, four were colonizers and six were not.

When considering abundance changes in stems ≥ 100 mm dbh, there was no indication that colonizers performed differently than non-colonizers. Excluding slope-specialists, 10 of 25 colonizing species decreased in abundance, while 24 of 53 non-colonizers declined. The four growth forms did not differ.

Swamp status. Swamp status was unrelated to population change. Considering stems ≥ 10 mm dbh, 26 of 51 swamp species declined in abundance (51%), whereas 78 of 167 non-swamp species declined (47%). For stems ≥ 100 mm dbh, 10 of 26 swamp species (38%) and 51 of 110 non-swamp (46%) declined. Neither difference, nor any for individual growth forms, reached statistical significance. Like colonizing species, though, swamp-specialists were over-represented among rapid increasers: seven of nine species increasing $\geq 5\%$ per year were swamp species, but only four of 18 species declining by $\geq 5\%$ per year were.

DISCUSSION

Nearly all shrubs and treelets that occur preferentially on the slopes of the 50 ha plot declined in abundance. We know the slopes around the side of the plateau are a wetter microhabitat during the dry season (Becker *et al.* 1988), and we assume that species more abundant there are less able to tolerate drought stress. We can support this assumption by casual observations on species distributions: some of the familiar slope-specialists at BCI – *Poulsenia armata*, *Olmedia aspera*, *Erythrina costaricana* and *Acalypha diversifolia* – are common along permanent streams in forests near BCI (there are no permanent streams on BCI). Further casual support comes from the genus *Piper*, which is particularly abundant in wet forests; five of its eight species in the 50 ha plot species are slope-specialists (all eight declined in abundance).

It seems certain that this group of shrubs and treelets that cannot tolerate long drought invaded the plateau forest at BCI during the wetter periods prior to 1966 but is now being eliminated by the increased severity of the dry season. It is possible that the extreme dry season of 1983 is solely responsible; alternatively, it may be a continuing problem caused by recurring severe dry seasons. We cannot distinguish between the two alternatives now, but future censuses will. If the only problem for drought-intolerant species was 1983, then populations should level off and perhaps even climb by 1995 or 2000, when the plot will be censused anew. We are certain, however, that dry season length and severity is the crucial edaphic variable affecting population success and limiting species' ranges at BCI (Wright 1992, Wright & van Schaik 1994). Reduction of rainfall during the wet season is probably inconsequential (at least for trees) since water is never limiting then.

Why have moisture-demanding trees of larger stature not suffered population declines as consistently as shrubs and treelets? We anticipated that they would, largely because of two prominent canopy trees and strong slope-specialists that suffered severe declines in abundance: *Poulsenia armata* and *Ocotea whitei*. But other slope-specialists in the canopy, such as *Calophyllum longifolium*, have healthy populations. We suggest the following hypothesis to account for this division and the general decline of shrubs and treelets. As adults, some trees, like *Calophyllum*, have longer roots than others, like *Poulsenia*, long enough to reach water below the slopes during the dry season, but not from the plateau (which is higher and thus further from the water table, see Wright & van Schaik 1994); *Poulsenia* thus suffered high mortality at all sizes during the 1983 El Niño drought, whereas *Calophyllum* did not. But both species have drought-sensitive seedlings, and are thus largely restricted to the wetter areas within the plot. Likewise, there are shrubs and treelets with drought-sensitive seedlings that are restricted to moist regions, but nearly all have short root systems as adults (Becker & Castillo 1990, Wright 1992) and suffer from long dry seasons at BCI. (Some shrubs have other drought-adaptations and are widespread in the plot.)

During moderate dry seasons, there is presumably enough water near the surface of the slopes for these drought-intolerant plants, and this allowed their spread into the plot prior to 1966, when dry seasons were less severe. The swamp may remain wet even during the most severe dry seasons, so that swamp-specialists can persist despite the drying trend.

Are the slope-specialists becoming extinct on BCI, or will they persist in locally wet sites? It appears not – all shrubs and treelets that declined throughout the plot declined on both the slopes and the plateau; in fact, most species had similar rates of change in both regions. Thus, it seems that there is a group of drought-intolerant species headed for extinction at BCI, at least 16 treelets and shrubs, and perhaps as many as 30–40 including the large, drought-sensitive trees like *Poulsenia* and *Ocotea whitei*. Howe (1990) has already predicted that *Virola surinamensis*, a large, slope-specialist tree, will go extinct on BCI due to its inability to tolerate long dry seasons. He based his conclusion on seedling survival data, not having seen the 1982–1990 population data which bears out his prediction – a decline from 300 to 239 stems.

Species which preferentially invade light gaps in the 50 ha plot – what we called colonizing species – had an almost bimodal distribution of population change. A few species had very rapid increases, but the rest did worse than average. Condit *et al.* (1992b) and Hubbell & Foster (1990a) have stated two different hypotheses about factors affecting the populations of gap-colonizers. One is that the more open canopy caused by drought-induced mortality in 1983 (Becker & Smith 1990) created more recruitment opportunities for species that demand light gaps, leading to a population burst in the 10 mm size class by 1985 or 1990. The other hypothesis is that the 50 ha plot is undergoing succession because the area just north of the plot (plus 2 ha within the plot) was cleared around the turn of the century but has matured since. Ruderal species abundant in the near-by farmland maintained high sink populations within the old forest because of the large number of seeds entering, but these populations are now declining. Both factors may in fact be at work at the same time. There are some colonizing species which can obviously maintain high populations within gaps of old forest (all seven of the rapidly rising species would be examples), but other species such as *Apeiba tibourbou* and *Schefflera morototoni* which are abundant only in large clearings may not persist in the old forest and are now in decline (indeed, the latter dropped out of the plot between 1982 and 1985).

An obvious concern with these conclusions is the method for identifying edaphic preferences of individual species. Some associations with topographic regions may be due to factors having nothing to do with moisture or light preference. Artefactual correlations weaken our power to detect effects of moisture preference, but the trends we did detect should be robust with respect to this error and ought to appear even stronger if species with accidental associations were segregated. We eventually hope to get physiological information on

some species to define independently those that are drought-tolerant (Mulkey *et al.* 1994, Wright & van Schaik 1994).

The mean rate of absolute population change in the plot was 2.25% per year, a 20% increase or 16% decrease after eight years. About 10% of the populations in the plot are changing >5% per year (a 49% increase or 33% decrease over eight years). These seem like substantial rates for trees, which ought to have rather lethargic population trajectories due to their long life spans and slow growth (Condit *et al.* 1992b, Hubbell & Foster 1990a). Are such changes typical for tropical forests, or is BCI unusual because of the climatic shift taking place? In a study of 50 ha of dry forest in India, several tree and shrub species underwent severe declines in abundance – in just three years – due to elephant herbivory (R. Sukumar, unpublished data). Other studies in the tropics have been on much smaller areas with irregular censuses, and are very difficult to compare. For example, Manokaran & Kochummen (1987) documented some abrupt declines and increases in a 34-year record of tropical forest in Malaysia: *Shorea parvifolia* declined from 26 to 16 individuals in 16 years, and *Dacryodes puberula* from 14 to four in 34 years, both consistent declines of about 5% per year, but both are based on small samples. The empirical issue of stability in tree populations and community composition of tropical forests must be resolved by more large datasets, and large-scale plots are now under way at 11 sites in Africa, Asia and America (Condit 1995). Results from these will settle the matter.

These plots will offer a baseline for assessing the impact of global climate change on tropical forests. Long-term changes in precipitation can have tremendous effects on forests (Foster 1982a,b, Hartshorn 1992), and if Phillips & Gentry (1994) are correct, CO₂-fertilization may be changing forest-wide dynamics. We see perhaps 10% of the species at BCI headed for extinction because of a 25-year decline in precipitation. Understanding and even anticipating climatic effects on tropical forests will be crucial for their long-range conservation.

ACKNOWLEDGEMENTS

The Smithsonian Tropical Research Institute in Panama provided generous logistical and financial support for the censuses, and we thank Ira Rubinoff for his long-term support of the project. We also thank the field workers who contributed to the censuses on BCI, more than 100 people from 10 countries, and R. Pérez and S. Loo de Lao for their persistent work maintaining the plot and its database. The project has been supported by grants from the National Science Foundation, the Smithsonian Scholarly Studies Program, the Smithsonian Tropical Research Institute, the John D. and Catherine T. MacArthur Foundation, the World Wildlife Fund, the Earthwatch Center for Field Studies, the Geraldine R. Doge Foundation and the Alton Jones Foundation. This article is a scientific contribution from the Center for Tropical Forest Science, which is supported by the John D. and Catherine T. MacArthur Foundation.

LITERATURE CITED

- BECKER, P. & CASTILLO, A. 1990. Root architecture of shrubs and saplings in the understorey of a tropical moist forest in lowland Panama. *Biotropica* 22:242–249.
- BECKER, P., RABENOLD, P. E., IDOL, J. R. & SMITH, A. P. 1988. Water potential gradients for gaps and slopes in a Panamanian tropical moist forest's dry season. *Journal of Tropical Ecology* 4:173–184.
- BECKER, P. & SMITH, A. P. 1990. Spatial autocorrelation of solar radiation in a tropical moist forest understorey. *Agricultural and Forest Meteorology* 52:373–379.
- BOTKIN, D. B. & NISBET, R. A. 1992. Projecting the effects of climate change on biological diversity in forests. Pp. 277–293 in Peters, R. L. & Lovejoy, T. E. (eds). *Global warming and biological diversity*. Yale University Press, New Haven.
- BRADLEY, R. S., DIAZ, H. F., EISCHEID, J. K., JONES, P. D., KELLY, P. M. & GOODESS, C. M. 1987. Precipitation fluctuations over northern hemisphere land areas since the mid-19th century. *Science* 237:171–175.
- BUSH, M. B. & COLINVAUX, P. A. 1990. A pollen record of a complete glacial cycle from lowland Panama. *Journal of Vegetation Science* 1:105–118.
- BUSH, M. B., COLINVAUX, P. A., WIEMANN, M. C., PIPERNO, D. R. & LIU, K. 1990. Late pleistocene temperature depression and vegetation change in Ecuadorian Amazonia. *Quaternary Research* 34:330–345.
- CONDIT, R. 1995. Research in large, long-term tropical forest plots. *Trends in Ecology and Evolution* 10:18–22.
- CONDIT, R., HUBBELL, S. P. & FOSTER, R. B. 1992a. Recruitment near conspecific adults and the maintenance of tree and shrub diversity in a neotropical forest. *American Naturalist* 140:261–286.
- CONDIT, R., HUBBELL, S. P. & FOSTER, R. B. 1992b. Stability and change of a neotropical moist forest over a decade. *Bioscience* 42:822–828.
- CONDIT, R., HUBBELL, S. P. & FOSTER, R. B. 1993a. Identifying fast-growing native trees from the neotropics using data from a large, permanent census plot. *Forest Ecology and Management* 62:123–143.
- CONDIT, R., HUBBELL, S. P. & FOSTER, R. B. 1993b. Mortality and growth of a commercial hardwood, 'El Cativo', *Prioria copaifera*, in Panama. *Forest Ecology and Management* 62:107–122.
- CONDIT, R., HUBBELL, S. P. & FOSTER, R. B. 1995. Mortality rates of 205 neotropical tree species and the responses to a severe drought. *Ecological Monographs* 65:419–439.
- CROAT, T. R. 1978. *Flora of Barro Colorado Island*. Stanford University Press, Stanford, California.
- DALE, V. H. & FRANKLIN, J. F. 1989. Potential effects of climate change on stand development in the Pacific Northwest. *Canadian Journal of Forestry Research* 19:1581–1590.
- D'ARCY, W. G. 1987. *Flora of Panama. Part 1: Introduction and checklist*. Missouri Botanical Garden, St Louis, Missouri, USA. 328 + xxx pp.
- DAVIS, M. B. 1981. Quaternary history and the stability of forest communities. Pp. 132–153 in West, D. C., Shugart, H. H. & Botkin, D. B. (eds). *Forest succession: concepts and applications*. Springer-Verlag, New York.
- DELCOURT, P. A. & DELCOURT, H. R. 1987. *Long term forest dynamics of the temperate zone: a case study of late quaternary forests in eastern North America*. Springer-Verlag, New York.
- DIAZ, H. F., BRADLEY, R. S. & EISCHEID, J. K. 1989. Precipitation fluctuations over global land areas since the late 1800's. *Journal of Geophysical Research* 94:1195–1210.
- FOSTER, R. B. 1982a. Famine on Barro Colorado Island. Pp. 201–212 in Leigh, E. G., Jr, Rand, S. A. & Windsor, D. M. (eds). *The ecology of a tropical forest: seasonal rhythms and long-term changes*. Smithsonian Institution Press, Washington, DC.
- FOSTER, R. B. 1982b. The seasonal rhythm of fruitfall on Barro Colorado Island. Pp. 151–172 in Leigh, E. G., Jr, Rand, S. A. & Windsor, D. M. (eds). *The ecology of a tropical forest: seasonal rhythms and long-term changes*. Smithsonian Institution Press, Washington, DC.
- FRANKLIN, J. F., SWANSON, F. J., HARMON, M. E., PERRY, D. A., SPIES, T. A., DALE, V. H., MCKEE, A., FERRELL, W. K., MEANS, J. E., GREGORY, S. V., LATTIN, J. D., SCHOWALTER, T. D. & LARSEN, D. 1992. Effects of global climatic change on forests in northwestern North America. Pp. 244–257 in Peters, R. L. & Lovejoy, T. E. (eds). *Global warming and biological diversity*. Yale University Press, New Haven.
- HAMILTON, A. C. & TAYLOR, D. 1991. History of climate and forests in tropical Africa during the last 8 million years. *Climate Change* 19:65–78.
- HARTSHORN, G. S. 1992. Possible effects of global warming on the biological diversity in tropical forests. Pp. 137–146 in Peters, R. L. & Lovejoy, T. E. (eds). *Global warming and biological diversity*. Yale University Press, New Haven.
- HOWE, H. F. 1990. Survival and growth of juvenile *Virola surinamensis* in Panama: effects of herbivory and canopy closure. *Journal of Tropical Ecology* 6:259–280.

- HUBBELL, S. P. & FOSTER, R. B. 1983. Diversity of canopy trees in a neotropical forest and implications for conservation. Pp. 25–41 in Sutton, S. L., Whitmore, T. C. & Chadwick, A. C. (eds). *Tropical rain forest: ecology and management*. Blackwell Scientific Publications, Oxford.
- HUBBELL, S. P. & FOSTER, R. B. 1986a. Commonness and rarity in a neotropical forest: implications for tropical tree conservation. Pp. 205–231 in Soulé, M. (ed.). *Conservation biology: the science of scarcity and diversity*. Sinauer Associates, Inc., Sunderland, Massachusetts.
- HUBBELL, S. P. & FOSTER, R. B. 1986b. Canopy gaps and the dynamics of a neotropical forest. Pp. 77–96 in Crawley M. J. (ed.). *Plant ecology*. Blackwell Scientific Publications, Oxford.
- HUBBELL, S. P. & FOSTER, R. B. 1987. The spatial context of regeneration in a neotropical forest. Pp. 395–412 in Crawley, M., Edwards, P. J. & Gray, A. (eds). *Colonization, succession, and stability*. Blackwell Scientific Publications, Oxford, England.
- HUBBELL, S. P. & FOSTER, R. B. 1990a. Structure, dynamics, and equilibrium status of old-growth forest on Barro Colorado Island. Pp. 522–541 in Gentry, A. (ed.). *Four Neotropical rain forests*. Yale University Press, New Haven.
- HUBBELL, S. P. & FOSTER, R. B. 1990b. The fate of juvenile trees in a neotropical forest: implications for the natural maintenance of tropical tree diversity. Pp. 325–349 in Hadley, M. & Bawa, K. S. (eds). *Reproductive ecology of tropical forest plants*. Man and the Biosphere Series, Vol. 7. UNESCO/IUBS, Paris, and Parthenon Publishing, Carnforth, UK.
- HUBBELL, S. P. & FOSTER, R. B. 1992. Short-term population dynamics of a neotropical forest: why ecological research matters to tropical conservation and management. *Oikos* 63:48–61.
- LEIGH, E. G., Jr, RAND, S. A. & WINDSOR, D. M. (eds). 1982. *The ecology of a tropical forest: seasonal rhythms and long-term changes*. Smithsonian Institution Press, Washington, DC.
- LEIGH, E. G., Jr, WINDSOR, D. M., RAND, S. A. & FOSTER, R. B. 1990. The impact of the 'El Niño' drought of 1982–1983 on a Panamanian semideciduous forest. Pp. 473–486 in Glynn, P. W. (ed.). *Global ecological consequences of the 1982–1983 El Niño–Southern Oscillation*. Elsevier Press.
- MANOKARAN, N. & KOCHUMMEN, K. M. 1987. Recruitment, growth and mortality of tree species in a lowland dipterocarp forest in Peninsular Malaysia. *Journal of Tropical Ecology* 3:315–330.
- MANOKARAN, N., LAFRANKIE, J. V., KOCHUMMEN, K. M., QUAH, E. S., KLAHN, J., ASHTON, P. S. & HUBBELL, S. P. 1992. *Stand table and distribution of species in the 50-ha research plot at Pasoh Forest Reserve*. Forest Research Institute of Malaysia, Research Data No. 1. 454 pp. Kepong, Malaysia.
- MULKEY, S. S., SMITH, A. P., WRIGHT, S. J., MACHADO, J. L. & DUDLEY, R. 1994. Contrasting leaf phenotypes control seasonal variation in water loss in a tropical forest shrub. *Proceedings of the National Academy of Sciences* 89:9084–9088.
- O'BRIEN, S. T., HAYDEN, B. P. & SHUGART, H. H. 1992. Global climatic change, hurricanes, and a tropical forest. *Climatic Change* 22:175–190.
- OVERPECK, J. T., RIND, D. & GOLDBERG, R. 1990. Climate-induced changes in forest disturbance and vegetation. *Nature* 343:51–53.
- PASTOR, J. & POST, W. M. 1988. Response of northern forests to CO₂-induced climate change. *Nature* 344:55–58.
- PHILLIPS, O. L. & GENTRY, A. H. 1994. Increasing turnover through time in tropical forests. *Science* 263:954–958.
- PHILLIPS, O. L., HALL, P., GENTRY, A. H., SAWYER, S. A. & VÁSQUEZ, R. 1994. Dynamics and species richness of tropical rain forests. *Proceedings of the National Academy of Science* 91:2805–2809.
- SHUGART, H. H. & SMITH, T. M. 1992. Using computer models to project ecosystem response, habitat change, and wildlife diversity. Pp. 147–157 in Peters, R. L. & Lovejoy, T. E. (eds). *Global warming and biological diversity*. Yale University Press, New Haven.
- SOLOMON, A. M. 1986. Transient response of forests to CO₂-induced climate change: simulation modeling experiments in eastern North America. *Oecologia* 68:567–579.
- SUKUMAR, R., DATTARAJA, H. S., SURESH, H. S., RADHAKRISHNAN, J., VASUDEVA, R., NIRMALA S. & JOSHI, N. V. 1992. Long-term monitoring of vegetation in a tropical deciduous forest in Mudumalai, southern India. *Current Science* 62:608–616.
- SUKUMAR, R., RAMESH, R., PANT, R. K. & RAJAGOPALAN, G. 1993. A δ¹³C record of late Quaternary climate change from tropical peats in southern India. *Nature* 364:703–706.
- SWAINE, M. D. & WHITMORE, T. C. 1988. On the definition of ecological species groups in tropical rain forests. *Vegetatio* 75:81–86.
- URBAN, D. L., HARMON, M. E. & HALPERN, C. B. 1993. Potential response of Pacific northwestern forests to climatic change, effects of stand and initial composition. *Climatic Change* 23:247–266.
- WELDEN, C. W., HEWETT, S. W., HUBBELL, S. P. & FOSTER, R. B. 1991. Survival, growth, and recruitment of saplings in canopy gaps and forest understorey on Barro Colorado Island, Panamá. *Ecology* 72:35–50.
- WINDSOR, D. M. 1990. *Climate and moisture variability in a tropical forest: long-term records from Barro Colorado Island, Panamá*. Smithsonian Contribution to the Earth Sciences, No. 29. Smithsonian Institution Press, Washington, DC.

- WINDSOR, D. M., RAND, A. S. & RAND, W. M. 1990. Características de la precipitación en la isla de Barro Colorado. Pp. 53–71 in Leigh, E. G., Rand, A. S. & Windsor, D. M. (eds). *Ecología de un bosque tropical: ciclos estacionales y cambios a largo plazo*. Smithsonian Tropical Research Institute, Balboa, República de Panamá.
- WOODS, P. 1989. Effects of logging, drought, and fire on structure and composition of tropical forests in Sabah, Malaysia. *Biotropica* 21:290–298.
- WRIGHT, S. J. 1992. Seasonal drought, soil fertility and the species diversity of tropical forest plant communities. *Trends in Ecology and Evolution* 7:260–263.
- WRIGHT, S. J. & VAN SCHAİK, C. P. 1994. Light and the phenology of tropical trees. *American Naturalist* 143:192–199.
- ZIMMERMAN, J. K., EVERHAM, E. M., III, WAIDE, R. B., LODGE, D. J., TAYLOR, C. M. & BROKAW, N. V. L. 1994. Responses of tree species to hurricane winds in subtropical wet forest in Puerto Rico: implications for tropical tree life histories. *Journal of Ecology* 82:911–922.

Accepted 18 March 1995

APPENDIX

The 48 species names that have been changed since the 50 ha plot was initiated in 1981 or which do not appear in Croat (1978). The current name is the one appearing in Table 1. The eight species listed as *sp. nov.* were newly discovered in the 50 ha plot and remain undescribed.

Current name	Name in D'Arcy (1987)	Name in Croat (1978)
<i>Appunia seibertii</i>	<i>Appunia seibertii</i>	not appearing
<i>Ardisia guianensis</i>	<i>Ardisia guianensis</i>	not appearing
<i>Brosimum guinense</i>	<i>Brosimum guinense</i>	not appearing
<i>Chamaedorea tepejilote</i>	<i>Chamaedorea tepejilote</i>	<i>Chamaedorea wenlandiana</i>
<i>Changuava schippii</i>	<i>Psidium anglohondrensis</i>	<i>Psidium anglohondrensis</i>
<i>Chrysochlamys eclipes</i>	<i>Tovomitopsis nicaraguensis</i>	<i>Tovomitopsis nicaraguensis</i>
<i>Chrysophyllum argenteum</i>	<i>Cynodendron panamense</i>	<i>Cynodendron panamense</i>
<i>Erythroxylum macrophyllum</i>	<i>Erythroxylum macrophyllum</i>	not appearing
<i>Garcinia intermedia</i>	<i>Garcinia intermedia</i>	<i>Rheedia edulis</i>
<i>Garcinia madruno</i>	<i>Garcinia madruno</i>	<i>Rheedia acuminata</i>
<i>Guarea grandifolia</i>	<i>Guarea grandifolia</i>	<i>Guarea multiflora</i>
<i>Guarea sp. nov.</i>	not appearing	not appearing
<i>Heisteria acuminata</i>	<i>Heisteria acuminata</i>	<i>Heisteria longipes</i>
<i>Hyeronima alcheornoides</i>	<i>Hyeronima laxiflora</i>	<i>Hyeronima laxiflora</i>
<i>Inga acuminata</i>	not appearing	not appearing
<i>Lonchocarpus latifolia</i>	<i>Lonchocarpus latifolia</i>	<i>Lonchocarpus pentaphyllum</i>
<i>Lopimia dasypetala</i>	<i>Lopimia dasypetala</i>	<i>Pavonia dasypetala</i>
<i>Maclura tinctoria</i>	<i>Chlorophora tinctoria</i>	not appearing
<i>Malmea sp. nov.</i>	not appearing	<i>Crematosperma sp.</i>
<i>Myrospermum frutescens</i>	<i>Myrospermum frutescens</i>	not appearing
<i>Nectandra purpurea</i>	<i>Nectandra purpurea</i>	<i>Nectandra purpurescens</i>
<i>Nectandra sp. nov. 1</i>	not appearing	not appearing
<i>Nectandra sp. nov. 3</i>	not appearing	not appearing
<i>Ocotea puberula</i>	<i>Ocotea puberula</i>	<i>Ocotea pyramidata</i>
<i>Ocotea whitei</i>	<i>Ocotea whitei</i>	<i>Ocotea skutchii</i>
<i>Oenocarpus mapoura</i>	<i>Oenocarpus mapoura</i>	<i>Oenocarpus panamanus</i>
<i>Ormosia amazonica</i>	<i>Ormosia amazonica</i>	not appearing
<i>Ormosia croatii</i>	<i>Ormosia coccinea</i>	<i>Ormosia coccinea</i>
<i>Phoebe cinnamomifolia</i>	<i>Phoebe cinnamomifolia</i>	<i>Phoebe mexicana</i>
<i>Pochota quinata</i>	<i>Bombacopsis quinata</i>	<i>Bombacopsis quinata</i>
<i>Pochota sessilis</i>	<i>Bombacopsis sessilis</i>	<i>Bombacopsis sessilis</i>
<i>Pourouma bicolor</i>	<i>Pourouma guianensis</i>	<i>Pourouma guianensis</i>
<i>Pouteria reticulata</i>	<i>Pouteria unilocularis</i>	<i>Pouteria unilocularis</i>
<i>Protium sp. nov.</i>	not appearing	not appearing

APPENDIX I. (cont.)

Current name	Name in D'Arcy (1987)	Name in Croat (1978)
<i>Psychotria graciflora</i>	<i>Psychotria graciflora</i>	not appearing
<i>Pterocarpus belizensis</i>	<i>Pterocarpus belizensis</i>	not appearing
<i>Sapium aucuparium</i>	<i>Sapium caudatum</i>	both (now considered synonyms)
<i>Sapium sp. nov.</i>	not appearing	not appearing
<i>Schefflera morototoni</i>	<i>Didymopanax morototoni</i>	<i>Didymopanax morototoni</i>
<i>Senna dariensis</i>	<i>Senna dariensis</i>	<i>Cassia fruticosa</i>
<i>Socratea exorrhiza</i>	<i>Socratea exorrhiza</i>	<i>Socratea durissima</i>
<i>Solanum steyemarkii</i>	<i>Solanum argenteum</i>	<i>Solanum argenteum</i>
<i>Terminalia oblonga</i>	<i>Terminalia oblonga</i>	<i>Terminalia chiriquensis</i>
<i>Trichilia pallida</i>	<i>Trichilia pallida</i>	<i>Trichilia montana</i>
<i>Trichilia tuberculata</i>	<i>Trichilia tuberculata</i>	<i>Trichilia cipo</i>
<i>Trichospermum galeottii</i>	<i>Trichospermum galeottii</i>	<i>Trichospermum mexicanum</i>
<i>Urera baccifera</i>	<i>Urera baccifera</i>	not appearing
<i>Virola sp. nov.</i>	not appearing	not appearing