# Abundance, Distribution, and Population Growth of the Northern Elephant Seal (*Mirounga angustirostris*) in the United States from 1991 to 2010

Mark S. Lowry,<sup>1</sup> Richard Condit,<sup>2</sup> Brian Hatfield,<sup>3</sup> Sarah G. Allen,<sup>4</sup> Ryan Berger,<sup>5</sup> Patricia A. Morris,<sup>6</sup> Burney J. Le Boeuf,<sup>7</sup> and J. Reiter<sup>6</sup>

<sup>1</sup>Marine Mammal and Turtle Division, Southwest Fisheries Science Center, National Marine Fisheries Service, NOAA, 8901 La Jolla Shores Drive, La Jolla, CA 92037, USA

E-mail: mark.lowry@noaa.gov

<sup>2</sup>Smithsonian Tropical Research Institute, Unit 9100, Box 0948, DPO AA 34002-0948, USA,

and National Center for Ecological Analysis and Synthesis, 735 State Street, Santa Barbara, CA 93101, USA

<sup>3</sup>U.S. Geological Survey, Western Ecological Research Center, Highway I, PO Box 70, San Simeon, CA 93452, USA

<sup>4</sup>National Park Service, Point Reyes National Seashore, One Bear Valley Road, Point Reyes Station, CA 94956, USA

<sup>5</sup>Point Blue Conservation Science, 3820 Cypress Drive, #11, Petaluma, CA 94954, USA

Institute for Marine Science, University of California, Santa Cruz, CA 95064, USA

Chancellor's Office, 275 Clark Kerr Hall, University of California, Santa Cruz, CA 95064, USA

# Abstract

We report on the distribution and abundance of the northern elephant seal (Mirounga angustirostris) in the United States from 1991 to 2010. Pup production (i.e., births) was the principal metric used to characterize abundance, distribution, and population growth of the U.S. population and of each rookery in the U.S. Birth estimates were obtained from the literature and estimated from recent counts of adult females or counts of pups made during ground and aerial photographic surveys conducted during the pupping-breeding season at all rookeries in central California and the Channel Islands in southern California. A total of 40,684 pups were estimated to have been born at 11 rookeries in the U.S. in 2010. The two most productive rookeries as of 2010 were San Miguel Island (16,208 pups) and San Nicolas Island (10,882 pups). The Piedras Blancas rookery was not established in 1991 but has since grown to be the fourth largest rookery as of 2010. Rookeries grew most rapidly initially, presumably due to high immigration rates; then increased moderately, eventually becoming stable; and some declined in size. Since 1988, the U.S. population has been growing at an average annual rate of 3.8%. The multiplicative factor needed to estimate total population size from pup production is estimated at 4.4. Total U.S. population size in 2010 was estimated at 179,000 individuals. Using conservative estimates for population growth of northern elephant seals in Mexico, we estimate that the total population in Mexico and the U.S. in 2010 was between 210,000 and 239,000 individuals.

Key Words: population growth, marine mammal census, pinniped, northern elephant seal, *Mirounga angustirostris* 

## Introduction

The northern elephant seal (Mirounga angustiros*tris*) was nearly driven to extinction during the 19th century by commercial sealers (Townsend, 1912; Bartholomew & Hubbs, 1960). A few northern elephant seals (estimated at fewer than 100) survived at Guadalupe Island, Mexico (Bartholomew & Hubbs, 1960; Hoelzel et al., 1993). Subsequent population growth and range expansion during the 20th century has been well-documented (see Stewart et al., 1994). In the early 21st century, northern elephant seals could be found at many island and mainland haulout sites in western Mexico and the United States (Figure 1). The present population shows little genetic diversity (Hoelzel et al., 1993) as a result of the genetic bottleneck due to the catastrophic population reduction in the 19th century (Bonnell & Selander, 1974).

As the northern elephant seal population grew, population expansion, distribution, and growth were observed at all rookeries in the U.S. and Mexico (Stewart et al., 1994). Stewart et al. (1994) estimated that 28,000 pups were born in Mexico and the U.S. in 1991 and that the total population (including northern elephant seals in Mexico and the U.S.) was 112,000 of which 75% resided in the U.S.

The annual number of births (i.e., pup production) is used to depict population distribution and to estimate population growth and abundance



Figure 1. Northern elephant seal (Mirounga angustirostris) rookeries in the U.S. and Mexico

because not all age/sex classes are present on land at the same time. Northern elephant seal births have been estimated from counts made by biologists on the ground (Stewart et al., 1994) and from aerial photographs (Lowry, 2002). Aerial photographic counts of northern elephant seals used in this report were found to be precise and accurate (Lowry et al., 1996). Herein, we (1) provide new or revised birth estimates for northern elephant seals at all U.S. rookeries through 2010 and include previously published estimates from Stewart et al. (1994), (2) estimate population growth at each U.S. rookery and for the U.S. population, (3) derive a new factor for converting birth estimates to population abundance, and (4) provide a population estimate for northern elephant seals in the U.S. in 2010.

## Methods

## Censuses

Censuses of northern elephant seal rookeries at the Channel Islands in southern California and central California from Point Conception to Point Reyes, where almost the entire U.S. population resides, were made by biologists on the ground and from aerial photographic surveys conducted during the winter pupping-breeding season (December to March). A minimum of one census of live pups was made each year near the end of the breeding season; counts of dead pups were included in most of these censuses. At some locations, additional censuses were conducted earlier in the breeding season to document the total number of adult females.

#### Ground Surveys

Ground surveys were made by one or more biologists over several days, depending on the size of the island or rookery. Northern elephant seals were tallied (with the aid of mechanical handcounters when necessary) while being observed by eye or through a hand-held binocular (refer to Stewart et al., 1994, for more details).

## Aerial Photographic Surveys

Aerial photographic surveys were conducted with a twin-engine, high-winged Partenavia PN68 or Partenavia PN68-Observer model aircraft flown at a ground speed of 185 km/h and at an altitude between 213 and 243 m. Multiple overlapping photographic passes were made over coastlines and beaches to ensure that all northern elephant seals on land were photographed (Lowry et al., 1996).

From 1988 to 2005, census photographs were taken with a 126-mm format Chicago Aerial Industries, Inc. KA-45A or KA-76 camera equipped with forward motion compensation and operated at a cycle rate that achieved 67% overlap between adjacent frames. In 2010, the northern elephant seals were photographed with a Canon EOS-1Ds Mark III, 20 megapixel digital camera attached to a mount having forward motion compensation and operated at a cycle rate that achieved 40% overlap between adjacent frames. The geographical position of each photograph was recorded by linking the camera (mounted vertically inside the belly of the aircraft) to a computer and Global Positioning System (GPS) receiver. A 152-mm-focal-length lens was used with the 126-mm format camera, and a 50- or 85-mm-focal-length lens was used with the Canon EOS-1Ds Mark III digital camera. Each camera was set at an aperture of f/5.6 with a shutter speed between 1/400 and 1/5,000 s. Three types of film were used in the 126-mm format camera: (1) Kodak Aerochrome MS Film 2448, a very fine-grained, medium-speed, color transparency film, was used during 1987-1999; (2) Aerochrome HS Film SO-359, a very fine-grained, high-speed, color transparency film, was used from 1997 to 2005; and (3) KODAK Aerochrome III MS Film 2427, a very fine-grained, medium-speed, color-reversal aerial film was used in 2005.

With color film, northern elephant seals were counted through a 7-70X zoom binocular microscope as the transparencies were illuminated on a light table. Images of animals were counted and marked on a clear acetate sheet with a different colored pen for each age/sex class category. Marks on the acetate were compared and verified with overlapping photographs. With digital photographs, animals were counted on computer monitors using Adobe Photoshop CS5 Extended software. Prior to counting the northern elephant seals, mosaics of coastline segments were created from multiple digital images using Adobe Photoshop CS5 because digital photographs were too small and covered too small an area. As the animals were counted, they were recorded in one of the following age/sex class categories: (1) live pups, (2) dead pups, (3) juveniles, (4) adult females, and (5) subadult and adult males.

#### Estimate of Births

Births were estimated for all northern elephant seal rookeries in southern and central California during the winter pupping-breeding season. Births were estimated from counts of live and dead pups and by either of two methods from an estimate of the number of adult females present at the rookery during the entire pupping-breeding season. Le Boeuf & Reiter (1988) determined that 97.5% of adult females present at the rookery during the entire breeding season gave birth to a pup. Therefore, the number of births can be estimated as 97.5% of adult females during the entire breeding season. However, not all adult females are present at the same time during the pupping-breeding seasons (i.e., adult females are an asynchronous population) and were found to be present for approximately 32 or 33 d during the  $2\frac{1}{2}$  to 3 mo breeding season (Le Boeuf & Laws, 1994). Two methods have been devised for estimating the total number of adult females at the rookery: (1) total number of adult females was assumed to be the count of adult females at peak time (usually a specifically determined date at the end of January) plus the count of adult females 32 or 33 d prior and after that date; and (2) the total numbers of adult females was estimated with an equation that calculates the total number of adult females in an asynchronous population from three or more censuses of adult females during the breeding season (Condit et al., 2007). At this time, it is unknown which method is best. Ground counts of adult females can easily be done using the 32 or 33 d method, but it may require more than one person to census a large rookery in a single day. Aerial photographic surveys can census several large rookeries in a single day but are difficult or impossible to carry out at 32 or 33 d intervals due to weather and aircraft availability. These two methods or direct counts of pups, or a combination of them, were used by the authors to estimate total number of births:

- Direct count of live pups at the end of the breeding season was used at Piedras Blancas rookery in 1992 and 1993.
- Direct count of live and dead pups at or near the end of the breeding season was used at rookeries located at Santa Barbara Island (1983-2010), Piedras Blancas (1994-2001), Cape San Martin/Gorda (1992-2010), Point Reyes (1981-1991), and at Año Nuevo (1961-1967).
- 3. Taking 97.5% of the estimate of total number of adult females counted during three censuses 32 or 33 d apart was used at Point Reyes (1993-2010) and Piedras Blancas (2002-2010).
- 4. Taking 97.5% of the estimate of total number of adult females using an equation that estimates total number of adult females in an asynchronous population from three or more counts of adult females was used at the Año Nuevo rookery during 1968 to 2010 (Le Boeuf et al., 2011) and at San Nicolas Island rookery in 2010.

- 5. Dead pups counted on aerial photographs taken during the mid to late February aerial census had a correction factor applied to them to compensate for dead pups washed off the beach or buried. The correction factor was based on six counts of adult females at San Nicolas Island in 2010 that was used in the equation that estimated total number of adult females and, subsequently, live births. The total number of live pups counted from the mid to late February 2010 aerial photo census was subtracted from that value to estimate total pup mortality. Dead pups counted from the 2010 mid to late February aerial census were then estimated to represent 26.6% of total pup mortality, the inverse of which yielded a multiplier (3.76) that converts the number of dead pups counted from aerial photographs to total pup mortality. The 3.76 multiplier was then applied to the count of all dead pups from all aerial photographic censuses taken at all Channel Islands colonies during mid to late February 1988 to 2010 censuses to estimate total pup mortality, which was then added to the total number of live pups counted in the mid to late February aerial censuses to estimate births.
- 6. The rookery at Santa Barbara Island experiences high pup mortality rate due to narrow beaches that are susceptible to high tides and large waves. For aerial photo surveys of this rookery, either the maximum count of pups or 97.5% of adult females counted from a mid to late February aerial census was used to estimate births.
- 7. The rookery at San Clemente Island is very small and is censused during the last week in January. Births were estimated from the count of live plus dead pups, or from 97.5% of adult females, whichever was greater.

#### Population Growth and Trends

Northern elephant seal birth estimates were used to estimate U.S. population growth rate and the growth rate of each rookery in central and southern California. The annual rate of increase ( $\lambda$ ) is calculated by the equation  $\lambda = e^r$ , where *r* is the slope of the linear regression from a series of natural logtransformed (ln) counts (Eberhardt & Simmons, 1992) obtained at the same time each year.

A posteriori examination of northern elephant seal birth estimates indicated non-linear growth of birth estimates over time for individual northern elephant seal rookeries and for regional summations of birth estimates (e.g., central California, Channel Islands, and U.S. populations). While some rookeries showed an increase in births, others showed an increase followed by a decrease in later years, or a slowing of the growth rate as time progressed. Breakpoint linear regression analysis (also known as piecewise regression analysis) of natural log-transformed birth estimates identified breakpoints in time series data. The program *SegReg* (www.waterlog.info/segreg.htm) was used to determine the location of the breakpoints in the time series of birth estimates and to perform linear regression analysis (y = a + bx) on each segment timeinterval from which the growth rate could be calculated for each segment. In some cases, no breakpoints were detected in the time interval. For each time-interval segment, the following was tabulated: (1) the average annual rate of increase ( $\lambda$ ) calculated from the slope of the regression, (2) number of observations (y) having birth estimates used in the analysis, (3) coefficient of determination (R<sup>2</sup>), and (4) coefficient of variation (CV) of the average annual growth rate (slope of the regression divided by the standard error of the slope).

# Total Population Size from Birth Estimates

A multiplicative factor for estimating the total population size from birth estimates was estimated by constructing a large number of life tables based on published ranges of survival and fecundity rates of elephant seals. For any complete life table (fecundity plus gender-specific survival at every age), there is a corresponding stable population growth rate,  $\lambda$ , defined as  $N_t/N_{t-1}$ (population sizes in consecutive years) and stable age distribution, defined as  $n_x/N_t$ , where  $n_x$  is the population size at age x at any time, and N is the total population at the same time. To find  $\lambda$  and the age distribution of both sexes, we simulated  $n_x$  in a population subject to a specified life table until they stabilized. The pup multiplier was then found as  $M = N/n_0$ , where  $n_0$  is the number of pups born. Moreover, assuming half of all births are male, the sex ratio at every age also stabilizes, and a multiplier for both males and females can be found. Our goal was to replicate a large number of life tables, covering published ranges of survival and fecundity, in order to estimate M, its variance, and how it relates to  $\lambda$ .

To generate many life tables, demographic rates were collected from published studies on annual survival and fecundity in northern and southern (Mirounga leonina) elephant seals (Cooper & Stewart, 1983; Le Boeuf & Reiter, 1988; Hindell, 1991; Huber et al., 1991; Reiter & Le Boeuf, 1991; Clinton & Le Boeuf, 1993; Le Boeuf et al., 1994; Pistorius & Bester, 2002; McMahon et al., 2003; Pistorius et al., 2004; Condit et al., 2013). From each study, if annual rates were included from > 1 y, we averaged across years, but different colonies were kept separate. Survival rates were averaged separately in four juvenile ages and two broad age categories in adults (Table 1). Fecundity, defined as pups weaned per female (thus including pregnancy less pup mortality), was averaged in two age categories (Table 1). The highest and lowest rates found across all studies are shown in Table 1. These ranges were converted to logit-normal distributions that produced the matching

95% confidence interval—that is, we found the mean and standard deviation (SD) of a logit-normal distribution that would produce rates falling in each desired range (Table 1). Life tables were then constructed stochastically using random draws from these logitnormal distributions, assuming no correlation among any of the rates. We created 100,000 life tables and simulated each to find its corresponding  $\lambda$  and M. The results were assembled into seven bins based on population growth, and in each  $\lambda$  bin, the mean and SD of M were calculated. We report the mean  $\pm 1.96$  SD.

## Results

In 2010, 40,684 northern elephant seal pups are estimated to have been born in the U.S. (Table 2). Since 1988, the U.S. population has been growing (Figure 2) at an average annual rate of 3.8% ( $\lambda = 1,038$ ; Table 3). The largest northern elephant seal rookery in 2010 was located at San Miguel Island (16,208 births), followed by San Nicolas Island (10,882 births), Santa Rosa Island (5,946 births), and Piedras Blancas (4,469 births) (Table 2). Of these four rookeries, in 2010, the Santa Rosa Island

**Table 1.** Annual survival and fecundity rates gleaned from the literature on northern and southern elephant seals (*Mirounga angustirostris* and *M. leonina*); the logit mean and SD describe a Gaussian distribution of logit-transformed rates which match the range of rates taken from the literature.

Age (y)	Sex	Rate	Range	Logit mean ± SD
1	Both	Survival	0.33-0.78	$0.279 \pm 0.493$
2	Both	Survival	0.67-0.91	$1.511 \pm 0.401$
3	Both	Survival	0.74-0.82	$1.281 \pm 0.118$
4	Both	Survival	0.68-0.87	$1.327 \pm 0.287$
5-16	Female	Survival	0.78-0.89	$1.678 \pm 0.206$
17-21	Female	Survival	0.56-0.73	$0.618 \pm 0.188$
5-10	Male	Survival	0.67-0.74	$0.877 \pm 0.084$
10-15	Male	Survival	0.57-0.67	$0.495 \pm 0.107$
3	Female	Fecundity	0.13-0.36	$-1.238 \pm 0.331$
$\geq 4$	Female	Fecundity	0.72-0.94	$1.848 \pm 0.452$



Figure 2. Total number of northern elephant seal births at the Channel Islands in southern California, along the coast in central California, and in the U.S. as a whole

	U.S. total	132'				1,032				2,010				3,329					6,523			9,262	10,427	11,589	12,355	13,837	15,079
	Total	84				972				1,823				2,938					5,528			7,662	8,818	9,692	10,232	11,499	13,139
	San Clemente Island													$0^{a}$					1 <sup>a</sup>					4k	0	3ķ	2 <sup>k</sup>
Islands	Santa Barbara Island					11 <sup>a</sup>								$28^{\rm a}$					$74^{\rm a}$						25 <sup>k</sup>	$80^{k}$	93 <sup>k</sup>
Channel	San Nicolas Island	$48^{\mathrm{a}}$				83ª		$136^{a}$		171 <sup>a</sup>	$193^{a}$	$210^{a}$	354ª	$428^{a}$					693ª			$1,187^{*}$	$1,546^{\circ}$	2,312 <sup>w</sup>	$1,906^{a}$	$2,074^{a}$	$2,583^{a}$
	Santa Rosa Island																				0	$0^{a}$	$0^{\mathrm{a}}$	$0^{a}$	0ª	$0^{a}$	$\mathcal{Q}^{\mathrm{a}}$
	bnslel leugiM ns2	84ª				$878^{a}$				$1,624^{a}$				$2,482^{a}$	$3,088^{\circ}$		$3,547^{a}$	$4,014^{a}$	$4,760^{a}$	$5,447^{a}$	$5,531^{a}$	$6,475^{a}$	$7,272^{a}$	$7,376^{a}$	8,301ª	9,342ª	$10,459^{a}$
	Total	0	5	12 23	32	60	75	100	143	187	238	307	326	391	437	512	657	815	995	1,131	1,330	1,600	1,609	1,897	2,123	2,338	1,940
	Point Conception																										
nia	Piedras Blancas																										
tral Califor	Cape San Martin/ Gorda																						$3^{a}$	$18^{a}$	$7^{\mathrm{a}}$	$25^{a}$	$24^{a}$
Cer	bnslsI ovəuN oñA bnslnism bns		çı	$12^{\text{s}}$	$32^{g}$	$60^{\text{g}}$	75	$100^{g}$	$143^{g}$	$187^{\rm g}$	238	$307^{\text{B}}$	$326^{s}$	390⊧	$435^{g}$	495 <sup>s</sup>	622 <sup>g</sup>	755 <sup>g</sup>	891 <sup>g</sup>	±866	$1,146^{g}$	$1,368^{s}$	$1,313^{g}$	$1,511^{g}$	1,639	$1,854^{g}$	$1,541^{s}$
	South Farallon Islands													$1^{a}$	$2^{\mathrm{a}}$	$17^{a}$	$35^{a}$	$60^{a}$	$104^{a}$	$133^{a}$	$184^{a}$	$232^{a}$	292ª	$367^{a}$	475 <sup>a</sup>	$447^{a}$	$360^{a}$
	Point Reyes																						1 <sup>a</sup>	$1^{a}$	$2^{\rm a}$	$12^{a}$	$15^{a}$
	Year	1958 1959	1960	1962	1963	1964	1965	1966	1967	1968	1969	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985

Table 2. Birth estimates of northern elephant seals for rookeries in the United States

25

	Istot .2.U	14,690	15,009	16,902	19,229	19,804	21,407	23,057	23,054	25,332	25,895	25,923		27,609		29,196	29,648				37,378					40,684	al. (2011);	es); <sup>k</sup> Lowry	cause pups	- J J
	Internet	12,507	13,033	14,724	16,833	17,287	18,511	19,833	19,938	21,777	21,833	21,337		22,631		24,160	24,410				30,207					33,144	e Boeuf et	adult female	n Miguel be	0
	San Clemente Island	1 <sup>k</sup>	$1^{k}$	0	0 <sup>k</sup>	0 <sup>k</sup>	3ķ	З,	2	0	$12^{k}$	$16^k$	$18^{k}$	$17^{k}$	$13^{k}$	Ţk	$17^{k}$	20 <sup>i</sup>	19	24	46	29	46	63	57	571	r (2011); <sup>g</sup> L	hs = 0.975*	count at Sa	
Islands	Santa Barbara Island	56 <sup>k</sup>	75 <sup>k</sup>	69 <sup>k</sup>	$106^{k}$	69 <sup>k</sup>	45 <sup>k</sup>	$18^{k}$	$106^{k}$	47 <sup>k</sup>	110 <sup>k</sup>	112 <sup>k</sup>		34 <sup>k</sup>		$67^{k}$	50 <sup>k</sup>				55					51	)12); 'Berge	ups; or birt	to the 1958	
Channel	San Nicolas Island	$2,626^{a}$	2,835ª	$3,248^{k}$	4,925 <sup>k</sup>	$4,266^{k}$	4,739 <sup>k</sup>	5,775 <sup>k</sup>	$5,310^{k}$	$5,964^{k}$	$6,821^{k}$	$6,353^{k}$		$6,828^{k}$		$10,018^{k}$	$9,403^{k}$				$10,245^{\circ}$					$10,882^{i}$	en et al. (20	3.76*dead p	d was added	
	Santa Rosa Island	0ª	$\mathcal{Q}^{\mathrm{a}}$	$2^{\rm a}$	$3^{\rm a}$	$27^{\rm a}$	$86^{k}$	$67^{k}$	123 <sup>k</sup>	315 <sup>k</sup>	$186^{k}$	435 <sup>k</sup>		516 <sup>k</sup>		$1,527^{k}$	$1,575^{k}$				4,227					5,946	2007); *All	live pups +	icolas Islan	
	bnslzl ləngiM ns2	$9,824^{a}$	$10,120^{a}$	$11,405^{k}$	$11,799^{k}$	$12,929^{k}$	$13,638^{k}$	$13,970^{k}$	$14,397^{k}$	$15,451^{k}$	$14,704^{k}$	$14,421^{k}$		$15,236^{k}$		$12,541^{k}$	$13,365^{k}$				$15,634^{i}$					$16,208^{i}$	dams et al. (	ata ( $Births =$	t from San N	
																											ib. data; <sup>d</sup> A	v, unpub. da	e 1959 coun	
	IstoT	2,183	1,976	2,178	2,396	2,517	2,896	3,224	3,116	3,555	4,062	4,586	5,052	4,978	5,524	5,036	5,238	5,567	6,192	6,662	7,171	7,580	7,673	7,392	7,015	7,540	ervice, unpu	a; <sup>J</sup> M. Lowr	les): and the	same year.
	Point Conception																				6					õ	al Park Se	inpub. dat	idult fema	luring the
nia	Piedras Blancas							$1^i$	54	293	605	989	1,196	1,671	$1,912^{\circ}$	1,846	$1,941^{6}$	2,176	2,649	3,053	$3,514^{\circ}$	3,789	4,080	4,007	3,808	4,469	9); <sup>o</sup> Nationa	3. Hatfield, ı	bs = 0.975 * a	not visited d
ntral Califor	Cape San Martin/ Gorda	4ª	$13^{a}$	$13^{a}$	$67^{a}$	$65^{a}$	263ª	$371^{h}$	$416^{\rm h}$	471 <sup>i</sup>	277	511	506	122	$431^{i}$	224'	266	362	305	299	$231^{6}$	278	250	$184^{1}$	204	$101^{10}$	z Allen (199	Hatfield); H	ups: or birth	t both were
Cei	bnslal ovəuN oñA bnslnism bns	$1,726^{\circ}$	$1,548^{g}$	$1,738^{g}$	$1,905^{s}$	$2,004^{g}$	$2,200^{s}$	2,366 <sup>∉</sup>	$2,168^{g}$	$2,276^{s}$	$2,731^{g}$	$2,484^{g}$	$2,720^{s}$	$2,554^{g}$	$2,575^{g}$	$2,416^{s}$	$2,473^{g}$	$2,466^{g}$	$2,594^{s}$	$2,512^{g}$	$2,619^{e}$	$2,582^{s}$	2,470 <sup>g</sup>	$2,368^{\sharp}$	$2,199^{s}$	$2,144^{g}$	Sydeman &	vided by B.	8.76*dead p	n islands, bu
	South Farallon Islands	434ª	395ª	393ª	$380^{a}$	$396^{a}$	$352^{a}$	370	329'	287 <sup>r</sup>	299	312 <sup>r</sup>	274 <sup>r</sup>	250 <sup>r</sup>	198	174	156	136'	149'	156	158	$185^{6}$	$168^{c}$	$150^{6}$	124	120'	al. (1994); <sup>1</sup>	ib. data (pro	$\varepsilon + sana ani$	born at both
	Point Reyes	19ª	$20^{a}$	$34^{a}$	ъ 44 а	52ª	81 <sup>a</sup>	$116^{\circ}$	$149^{\text{b}}$	228 <sup>6</sup>	$150^{\circ}$	$290^{\circ}$	$356^{\circ}$	$381^{\text{b}}$	$408^{\circ}$	376°	$402^{\circ}$	$427^{\circ}$	495°	$642^{\circ}$	$643^{\circ}$	$746^{d}$	705∘	683°	$680^{\circ}$	698°	Stewart et a	neson, unpu	$(Births = l_i$	kely being
	Year	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	<sup>a</sup> From	<sup>h</sup> R. Jan	(2002)	were li

Lowry et al.



Figure 3. Northern elephant seal births at rookeries in the U.S.

rookery experienced the highest average annual growth rate (45.6%,  $\lambda = 1.456$ ), followed by San Nicolas Island (11.9%,  $\lambda = 1.119$ ) and Piedras Blancas (10.8%,  $\lambda = 1.108$ ). The rookeries at San Miguel Island appear to be leveling off ( $\lambda = 1.005$ ; Table 3). In 2010, the rookeries at the Farallon Islands, Año Nuevo, Cape San Martin/

Gorda, and Santa Barbara Island were declining (Figure 3; Table 3).

The multiplicative factor, M, needed to estimate total population size from the number of births was 4.4 when population growth rate  $\lambda$  was > 1.02 (Table 4), with a 95% confidence interval spanning 3.8 to 5.0. M did not vary much with  $\lambda$ , averaging nearly 4.2

**Table 3.** Average annual rate of increase ( $\lambda$ ), number of observations (n), coefficient of determination ( $\mathbb{R}^2$ ), and coefficient of variation ( $\mathbb{CV}$ ) for the rates of increase that were computed for year-intervals chosen by breakpoint regression analysis of log-transformed northern elephant seal birth estimates for rookeries in the U.S.

Rookery	Year interval	п	R <sup>2</sup>	λ	CV
Point Reyes	1981-1992	12	0.91	1.541	0.098
·	1993-2010	18	0.86	1.092	0.102
Farallon Islands	1973-1980	8	0.85	1.808	0.168
	1981-2010	30	0.84	0.955	-0.084
Año Nuevo	1961-1977	17	0.95	1.279	0.061
	1978-1995	18	0.89	1.048	0.088
	1996-2010	15	0.40	0.991	-0.337
Cape San Martin/Gorda	1981-1994	14	0.77	1.440	0.157
	1995-2010	16	0.29	0.951	-0.417
Piedras Blancas	1992-1995	4	0.89	8.090	0.251
	1996-2010	15	0.94	1.108	0.073
San Miguel Island	1958-1979	10	0.92	1.197	0.106
	1980-1990	11	0.95	1.067	0.076
	1991-2010	11	0.15	1.005	0.793
Santa Rosa Island	1985-2010	16	0.87	1.456	0.102
San Nicolas Island	1964-2010	27	0.92	1.119	0.059
Santa Barbara Island	1964-1983	5	0.30	1.051	0.872
	1983-2010	18	0.06	0.984	-1.015
San Clemente Island	1977-2010	24	0.87	1.155	0.081
Central California	1958-1970	11	0.93	1.547	0.092
	1971-1979	9	0.99	1.200	0.030
	1980-2010	31	0.96	1.059	0.036
Channel Islands	1958-1987	13	0.92	1.164	0.091
	1988-2010	14	0.95	1.034	0.065
United States	1958-1987	13	0.92	1.170	0.091
	1988-2010	14	0.96	1.038	0.058

**Table 4.** Estimated multipliers for deriving total population size from the number of northern elephant seal births (with 95% confidence limits [CL] in parentheses) as a function of population rate of increase ( $\lambda$ ; i.e., population growth rate); the multiplier (*M*) for females gives the total female population size from female pups, and likewise for males; *M* for the total population is their mean.

	Fer	nale	М	ale	Total population						
λ	Multiplier	95% CL	Multiplier	95% CL	Multiplier	95% CL					
0.95-0.96	4.70	(4.15-5.24)	3.62	(2.92-4.31)	4.16	(3.28-5.04)					
0.97-0.98	4.77	(4.27-5.28)	3.71	(3.08-4.34)	4.24	(3.43-5.05)					
0.99-1.00	4.83	(4.38-5.28)	3.78	(3.22-4.34)	4.31	(3.59-5.02)					
1.01-1.02	4.88	(4.47-5.28)	3.84	(3.35-4.33)	4.36	(3.72-4.99)					
1.02-1.03	4.89	(4.51-5.28)	3.86	(3.41-4.32)	4.38	(3.78-4.97)					
1.03-1.04	4.91	(4.57-5.25)	3.88	(3.48-4.28)	4.39	(3.87-4.92)					
≥ 1.04	4.91	(4.60-5.22)	3.88	(3.51-4.26)	4.40	(3.91-4.88)					



Figure 4. Percentage of U.S. northern elephant seal births recorded at rookeries at the Channel Islands in southern California and along the coast in central California

at the lowest rates of population growth and never higher than 4.4 (Table 4). Rather, most uncertainty in M was due to uncertainty in demographic rates (Table 1). With fecundity (including pre-weaning pup survival) as low as it has been observed (forcing survival to be its highest), M reached values near 5.0. Conversely, if fecundity were as high as it has been observed (so that survival must be low), then Mfell < 4.0. The pup multiplier for males (male pups converted to total males) was lower than for females (Table 4) because of lower survival rates in adult males relative to females (Table 1).

Total U.S. northern elephant seal population size in 2010 is estimated at 179,000 (2.5% = 159,000, 97.5% = 199,000). Expansion of the U.S. population of northern elephant seals was due primarily to growth at Channel Islands rookeries in southern California (Figure 2). Channel Islands rookeries accounted for 81.5% of births in 2010, with the remaining 18.5% of births occurring in central California (Figure 4). Since 1988, northern elephant seal births at the Channel Islands have been increasing at an average annual growth rate of 3.4% ( $\lambda$  = 1.034, CV = 0.065), and at an average annual growth rate of 5.9% ( $\lambda$  = 1.059, CV = 0.036; Table 3) in central California since 1980.

#### Discussion

The U.S. population estimate for 2010 exceeds the last estimate made in 1991 by Stewart et al. (1994). Although we do not have current birth estimate data for the population of northern elephant seals in Mexico, if both populations experienced similar growth rates since the 1991 estimate, then the total northern elephant seal population in 2010 was roughly 239,000 individuals. However, Stewart et al. reported that the population in Mexico had stabilized in 1991. If the population in Mexico did not grow, then the total population estimate for the species in 2010 would be roughly 210,000 individuals. It would be reasonable to say, then, that in 2010 the combined northern elephant seal population in Mexico and the U.S. included 210,000 to 239,000 individuals.

The U.S. northern elephant seal population continues to grow, with most of the growth occurring at the larger rookeries in southern California. New rookeries show a period of extremely fast growth in the first few years, presumably due to immigration because the initial rate of growth is not biologically feasible, followed by a gradual slowing of the growth rate. Eventually, some rookeries, such as those at the Farallon Islands, Año Nuevo Point and Island, Cape San Martin/Gorda, and Santa Barbara Island, experienced negative growth. As the Farallon Islands rookery diminished in size, the Point Reyes rookery increase probably resulted from northern elephant seals moving there from the Farallon Islands, Piedras Blancas, and Año Nuevo which are nearby (Condit et al., 2013). Growth at Piedras Blancas occurred as the Año Nuevo rookery began to decrease and as the San Miguel Island rookery decreased in 2000 and 2001. The Santa Rosa Island rookery increased due to immigration from San Miguel Island (which is next to it) as is shown by decreases in births at San Miguel Island in 2000 and 2001. Eventually, the number of births at San Miguel Island increased such that by 2005 and 2010, the number of births at that rookery exceeded the previous high level. When that happened, there were enough northern elephant seals at Santa Rosa Island to continue population expansion at that island. On occasion, a few pups have been produced at Castle Rock (Point Saint George) in northern California near Crescent City (Hodder et al., 1998). We did not count that site in 2010, but it generally has produced no more than 11 pups. Still further north, elephant seals have bred at Shell Island, Oregon, and Race Rocks and other sites in the Strait of Juan de Fuca, Washington, and British Columbia, but only a few pups are produced at these sites, and they rarely survive to weaning because of inclement weather (Hodder et al., 1998; Jeffries et al., 2000).

The pup multiplier we present is similar to those for earlier estimates for gray seals (*Halichoerus grypus*) and southern elephant seals, but no prior estimates have separated males and females or produced estimates of uncertainty. The uncertainty in *M*, caused mostly by uncertainty in demographic rates, adds about 20% uncertainty in total U.S. population size, thus our population estimate spans 159,000 to 199,000.

Northern elephant seal population expansion is expected to continue but at a reduced rate due to increased pup mortality at crowded rookeries. Sandy beaches (preferred terrestrial habitat) at San Nicolas Island and San Miguel Island are all almost totally occupied by elephant seals during the pupping-breeding season. The same will happen soon at Santa Rosa Island. Santa Cruz Island, Anacapa Island, and other Channel Islands either have a limited number of sandy beaches or none at all. The southern California coastline has suitable beaches, but human presence will probably keep northern elephant seals from occupying them. Point Reyes and Piedras Blancas rookeries have suitable and ample beach space for rookery growth nearby. The Point Conception rookery is extremely small as it lies in a small cove beneath a towering cliff and is occupied by Pacific harbor seals (*Phoca vitulina richardii*). There is, however, ample beach space available north of Point Arguello within Vandenberg Air Force Base that could possibly be occupied by northern elephant seals in the future.

## Acknowledgments

We thank many colleagues, students, and volunteers for years of censuses of northern elephant seals. There are too many of them to list, but they know who they are, and their contributions are included in this report. H. Huber, W. Sydeman, and R. Bradley provided long-term support at the Farallon Islands. D. Adams, D. Roberts, and D. Press provided longterm support at Point Reyes. K. Karako assisted with many elephant seal counts at the Piedras Blancas rookery. W. Perryman of the Southwest Fisheries Science Center (SWFSC) provided photographic equipment for aerial photographic surveys. Aircraft for conducting aerial photographic surveys were chartered from Everingham Brothers (1988-1990) and Aspen Helicopters (1991-2010). Vessels for transport to Santa Barbara Island were provided by Channel Islands National Park or chartered from Peter Howorth. The U.S. Navy assisted with surveys at San Clemente Island and San Nicolas Island. The U.S. Air Force assisted with surveys at Vandenberg Air Force Base. Some of the research was supported in part by several grants from the National Science Foundation, and funding from the National Oceanic and Atmospheric Administration and the National Park Service. The surveys conducted on the Farallon Islands and Point Reyes were covered under National Marine Fisheries Service permit #373-1575; and Año Nuevo censuses were covered under permit #87-1743-04. All SWFSC surveys were conducted under Marine Mammal Research Permits 347, 404, 684, 704, 774-1437, 774-1714, and 14097; and National Marine Sanctuary Permits GFNMS/MBNMS/ CINMS-04-98, MULTI-2002-003, MULTI-2003-003, and MULTI-2008-003. All authors were involved in the collection of data and approved the final manuscript. The report was improved by comments provided by J. Barlow, W. Perryman, J. Laake, W. Perrin, and two anonymous reviewers.

#### Literature Cited

- Adams, D., Jensen, H., Nevins, H., Truchinski, K., Roberts, D., & Allen, S. (2007). Northern elephant seal monitoring 2005-2007 report, Point Reyes National Seashore (Natural Resource Technical Report NPS/SFAN/NRTR–2008/085). Fort Collins, CO: National Park Service. 30 pp.
- Allen, S., Codde, S., Jensen, H., Wheeler, E., Becker, B., Press, D., & Roberts, D. (2012). Northern elephant seal monitoring (Mirounga angustirostris) at Point Reyes National Seashore: 2008-2009 breeding seasons (Natural Resource Technical Report NPS/SFAN/NRTR— 2012/649). Fort Collins, CO: National Park Service. 28 pp.
- Bartholomew, G. A., & Hubbs, C. L. (1960). Population growth and seasonal movements of the northern elephant seal, *Mirounga angustirostris. Mammalia*, 24, 313-324.

- Berger, R. W. (2011). Population size and reproductive performance of northern elephant seals on the South Farallon Islands, 2010-2011 (PRBO Contribution Number 1803). Unpublished report to the U.S. Fish and Wildlife Service, PRBO Conservation Science, Petaluma, CA. 21 pp.
- Bonnell, M. L., & Selander, R. K. (1974). Elephant seals: Genetic variation and near extinction. *Science*, 184, 908-909. http://dx.doi.org/10.1126/science.184.4139.908
- Clinton, W. L., & Le Boeuf, B. J. (1993). Sexual selection's effects on male life history and the pattern of male mortality. *Ecology*, 74, 1884-1892. http://dx.doi. org/10.2307/1939945
- Condit, R., Le Boeuf, B. J., Morris, P. A., & Sylvan, M. (2007). Estimating population size in asynchronous aggregations: A model and test with elephant seal counts. *Marine Mammal Science*, 23, 834-855. http:// dx.doi.org/10.1111/j.1748-7692.2007.00141.x
- Condit, R., Reiter, J., Morris, P. A., Berger, R., Allen, S. G., & Le Boeuf, B. J. (2013). Lifetime survival rates and senescence in northern elephant seals. *Marine Mammal Science*, early view. http://dx.doi.org/10.1111/ mms.12025
- Cooper, C. F., & Stewart, B. S. (1983). Demography of northern elephant seals, 1911-1982. *Science*, 219, 969-971. http://dx.doi.org/10.1126/science.219.4587.969
- Eberhardt, L. L., & Simmons, M. A. (1992). Assessing rates of increase from trend data. *Journal of Wildlife Management*, 56, 603-610. http://dx.doi.org/10.2307/3808878
- Hindell, M. A. (1991). Some life-history parameters of a declining population of southern elephant seals, *Mirounga leonina. Journal of Animal Ecology*, 60, 119-134. http://dx.doi.org/10.2307/5449
- Hodder, J., Brown, R., & Cziela, C. (1998). The northern elephant seal in Oregon: A pupping range extension and onshore occurrence. *Marine Mammal Science*, 14, 873-881. http://dx.doi.org/10.1111/j.1748-7692.1998.tb00772.x
- Hoelzel, A. R., Halley, J., O'Brien, S. J., Campagna, C., Arnbom, T., Le Boeuf, B. J., . . . Dover, G. A. (1993). Elephant seal genetic variation and the use of simulation models to investigate historical population bottlenecks. *Journal of Heredity*, 84, 443-449.
- Huber, H. R., Rovetta, A. C., Fry, L. A., & Johnston, S. (1991). Age-specific natality of northern elephant seals at the South Farallon Islands, California. *Journal of Mammalogy*, 72, 525-534. http://dx.doi. org/10.2307/1382136
- Jeffries, S. J., Gearin, P. J., Huber, H. R., Saul, D. L., & Pruett, D. A. (2000). Atlas of seal and sea lion haulout sites in Washington. Washington Department of Fish and Wildlife, Wildlife Science Division, 600 Capitol Way North, Olympia, WA 98501.
- Le Boeuf, B. J., & Laws, R. M. (1994). Elephant seals: An introduction to the genus. In B. J. Le Boeuf & R. M. Laws (Eds.), *Elephant seals: Population ecology, behavior, and physiology* (pp. 1-28). Berkeley: University of California Press.
- Le Boeuf, B. J., & Reiter, J. (1988). Lifetime reproductive success in northern elephant seals. In T. H. Clutton-

Brock (Ed.), *Reproductive success: Studies of individual* variation in contrasting breeding systems (pp. 344-362). Chicago and London: The University of Chicago Press.

- Le Boeuf, B. J., Morris, P., & Reiter, J. (1994). Juvenile survivorship of northern elephant seals from Año Nuevo. In B. J. Le Boeuf & R. M. Laws (Eds.), *Elephant seals: Population ecology, behavior, and physiology* (pp. 121-136). Berkeley: University of California Press.
- Le Boeuf, B. J., Condit, R., Morris, P. A., & Reiter, J. (2011). The northern elephant seal (*Mirounga angustiros-tris*) rookery at Año Nuevo: A case study in colonization. Aquatic Mammals, 37(4), 486-501. http://dx.doi. org/10.1578/AM.37.4.2011.486
- Lowry, M. S. (2002). Counts of northern elephant seals at rookeries in the Southern California Bight: 1981-2001 (NOAA Technical Memorandum NMFS NOAA-TM-NMFS-SWFSC-345). Washington, DC: National Oceanic and Atmospheric Administration.
- Lowry, M. S., Perryman, W. L., Lynn, M. S., Westlake, R. L., & Julian, F. (1996). Counts of northern elephant seals (*Mirounga angustirostris*) from large-format aerial photographs taken at rookeries in southern California during the breeding season. U.S. Fishery Bulletin, 94, 176-185.
- McMahon, C. R., Burton, H. R., & Bester, M. N. (2003). A demographic comparison of two southern elephant seal populations. *Journal of Animal Ecology*, 72, 61-74. http://dx.doi.org/10.1046/j.1365-2656.2003.00685.x
- Pistorius, P. A., & Bester, M. N. (2002). A longitudinal study of senescence in a pinniped. *Canadian Journal* of Zoology, 80, 395-401. http://dx.doi.org/10.1139/z02-017
- Pistorius, P. A., Bester, M. N., Lewis, M. N., Taylor, F. E., Campagna, C., & Kirkman, S. P. (2004). Adult female survival, population trend, and the implications of early primiparity in a capital breeder, the southern elephant seal (*Mirounga leonina*). Journal of Zoology, 263, 107. http://dx.doi.org/10.1017/S0952836904004984
- Reiter, J., & Le Boeuf, B. J. (1991). Life history consequences of variation in age at primiparity in northern elephant seals. *Behavioral Ecology and Sociobiology*, 28, 153-160. http://dx.doi.org/10.1007/BF00172166
- Stewart, B. S., Yochem, P. K., Huber, H. R., DeLong, R. L., Jameson, R., Sydeman, W., . . . Le Boeuf, B. J. (1994). Population recovery and status of the northern elephant seal, *Mirounga angustirostris*. In B. J. Le Boeuf & R. M. Laws (Eds.), *Elephant seals: Population ecology, behavior, and physiology* (pp. 29-48). Berkeley: University of California Press.
- Sydeman, W. J., & Allen, S. G. (1999). Pinniped population dynamics in central California: Correlations with sea surface temperature and upwelling indices. *Marine Mammal Science*, 15, 446-461. http://dx.doi. org/10.1111/j.1748-7692.1999.tb00812.x
- Townsend, C. H. (1912). The northern elephant seal Macrorhinus angustirostris (Gill). Zoologica, 1, 159-173.