

ORIGINAL ARTICLE

Viability of small seeds found in feces of the Central American tapir on Barro Colorado Island, Panama

Paula I. CAPECE¹, Enzo ALIAGA-ROSSEL^{1,2} and Patrick A. JANSEN^{3,4}

¹Ecology, Evolution and Conservation Biology Program, University of Hawaii at Manoa, Honolulu, HI, USA, ²Institute of Ecology, Higher University of San Andres, La Paz, Bolivia, ³Smithsonian Tropical Research Institute, Ancón, Panama City, Panama and ⁴Center for Ecosystem Studies, Wageningen University, Wageningen, Netherlands

Abstract

Tapirs are known as effective dispersers of large-seeded tree species, but their role in dispersing small-seeded plant species has yet to be established. Tapir feces have been reported to contain large numbers of small seeds, but whether these are viable has rarely been evaluated. We determined the abundance and viability of small seeds in feces of Central American tapir (*Tapirus bairdii*) on Barro Colorado Island, Panama. A total of 72 fecal samples were collected opportunistically from 4 tapir latrine sites. Seeds were manually extracted from feces and classified by size. Seed viability was estimated by opening each seed and examining for the presence of at least 1 intact firm white endosperm. In total, we obtained 8166 seeds of at least 16 plant species. Small-seeded species dominated, with 96% of all seeds found measuring <5 mm. The canopy tree *Laetia procera* was the most abundant species in the samples. Of all small seeds found, 69% contained an intact endosperm and appeared viable. This suggests that small seeds, like large seeds, often pass through the digestive tract of *T. bairdii* intact. Thus, tapirs potentially serve as effective dispersers of a wide range of small-seeded plant species.

Key words: Central America, megafauna, seed dispersal, seed size, tapir

INTRODUCTION

Seed dispersal by tapirs (*Tapirus* spp.) is an important determinant of the spatial distribution of plants in Neotropical forests (Forget & Sabatier 1997; Fragoso *et al.* 2003). Tapirs are the largest extant terrestrial mam-

mals in Neotropical forests, are able to swallow large seeds and can ingest a large quantity of seeds while feeding (Terwilliger 1978; Naranjo 1995). Tapirs possess a 'primitive' digestive tract characterized by incomplete digestion; therefore, many seeds can pass through the tapir gut intact (Janzen 1984; Bodmer 1991). Long seed retention times in the gut (Janzen 1981) and large home-range sizes (Foerster & Vaughan 2002) contribute to the tapir's ability to disperse seeds over long distances, in some cases to suitable sites beyond the range of seed-consuming larvae (Fragoso *et al.* 2003).

Correspondence: Paula I. Capece, National Park Service, 1978 Island Ford Parkway, Sandy Springs, GA 30350, USA.
Email: paula_capece@nps.gov

Tapirs are considered effective dispersers of plant species with large, thick-walled seeds; that is, they tend to deposit seeds unharmed to suitable germination sites away from the parent plant (see Schupp 1993 and Jordano & Schupp 2000 for a detailed discussion of effective dispersal). For example, *Tapirus terrestris* (Linnaeus, 1758) is an effective disperser of large-seeded palms (Rodrigues *et al.* 1993; Fragoso & Huffman 2000; Quiroga-Castro & Roldan 2001; Tobler *et al.* 2010) and *Tapirus bairdii* (Gill, 1865) is an effective disperser of large-seeded tree species, including *Caesalpinia coriaria* (Janzen 1982), *Guazuma ulmifolia* (Janzen 1982) and *Manilkara zapota* (O'Farrill *et al.* 2006, 2012). However, tapirs also feed on fruits that contain many small seeds and are known to ingest small seeds while browsing (Janzen 1984). Fragoso and Huffman (2000) report that seeds of 4 small-seeded species germinated after they were extracted from *T. terrestris* feces. Small seeds have been found in *T. terrestris* stomach contents (Henry *et al.* 2000) and feces (Tófoli 2006; Zorzi 2009; Tobler *et al.* 2010), as well as in *T. bairdii* feces (Naranjo 1995). Whether tapirs are effective dispersers of these small seeds is partially determined by the degree to which small seeds survive passage through the digestive tract intact, which is largely unknown.

We examined the potential of the Central American tapir (*T. bairdii*) as a disperser of small-seeded plant species by estimating the viability of seeds found in tapir feces on Barro Colorado Island, Panama. We also recorded the abundance and diversity of small seeds in feces.

MATERIALS AND METHODS

Study site and species

Barro Colorado Island (BCI), Panama (9°9'N, 79°51'W) is a 16 km² island covered with tropical lowland moist forest. BCI has an average temperature of 27 °C, an average yearly rainfall of 2600 mm, a rainy season from May through December and a marked dry season from January through April. The island has a heterogeneous landscape with numerous creeks, most of which are seasonally dry (Leigh *et al.* 1996).

Tapirus bairdii is 1 of 4 species of the family Tapiridae of the order Perissodactyla and the only remaining Pleistocene megafauna in Central America (Janzen 1982). It is distributed intermittently from southern Mexico to northwestern Colombia in humid habitats

from sea level to 3600 m (IUCN 2012). The diet of *T. bairdii* consists of a variety of plant species and plant parts, including leaves, stems and fruits (Terwilliger 1978; Naranjo 1995; Olmos 1997; Tobler 2002). *T. bairdii* is more active during nocturnal hours (Foerster & Vaughn 2002). It is known to use latrines or sites of repeated defecation (Flesher 1999; Naranjo 2009).

The diet of *T. bairdii* on BCI has been studied by Terwilliger (1978) through direct observation of the feeding behavior of individual tapirs. To our knowledge, no studies have examined fecal samples on BCI. Estimates of the tapir population on BCI have varied greatly over time (Enders 1935; Enders 1939; Eisenberg & Thorington 1973; Terwilliger 1978; Eisenberg 1980; Glanz 1996). More recent estimates of *T. bairdii* density on BCI obtained from diurnal and nocturnal census efforts are 0.19 and 0.36 individuals/km², respectively (Wright *et al.* 1994). The current population size of tapirs on BCI is unknown (R. W. Kays, pers. comm.).

Sampling

We collected fresh fecal samples (dung boluses <1 week old as judged by inspection of odor, color and integrity of fecal material) at 4 tapir latrine locations along the northern shore of BCI, near the terminal portions of the 'Standley' and 'Miller' trails, during May–Aug 2003. All 4 latrines were located within a straight-line distance of approximately 1.5 km (approximately 3 km ground distance), with 2 of the latrines in close proximity (approximately 50 m). Fecal samples were slowly dried in a dry closet for 3 days at approximately 35 °C and then spread out on trays in a dark air-conditioned laboratory for an additional 7 days until completely dry. Each fecal sample was packaged individually in a paper bag and assigned a number for identification, then stored in plastic with desiccant until processing, which was completed within 8 weeks. We extracted the seeds from dry feces manually. This proved more efficient than seed extraction using different-sized wire mesh screens (as in Naranjo 1995; Fragoso & Huffman 2000), because many small seeds passed directly through 5 mm² mesh sieves and could not be effectively separated from browse material with smaller-mesh sieves. We classified seeds as small (S < 5 mm), medium (M = 5–20 mm) or large (L > 20 mm) and identified seeds to 'morphospecies'. We attempted to identify morphospecies to species through comparison with BCI herbarium samples and with the assistance of Osvaldo Calderón and Andres Hernandez (BCI botanists). Viability was estimat-

ed by the cut test (Terry *et al.* 2003); opening each unimbibed seed and examining under magnification for the presence of at least 1 intact firm white endosperm (as in Fragoso & Huffman 2000). Although the cut test method is indirect and may vary by species, it has been used successfully to determine viability for tapir ingested seeds (Fragoso & Huffman 2000), for small seeds dispersed by frugivorous tortoises (Strong & Fragoso 2006) and for other small seeds (*Piper* spp.) where germination tests failed to distinguish between viability and dormancy (Daws *et al.* 2002).

RESULTS

Of 72 fecal samples obtained, 52 from 3 tapir latrine locations in the early wet season comprised approximately 3900 cm³ of fecal matter containing 1084 seeds (Table 1). A total of 20 samples from a fourth latrine location during the mid-wet season comprised approximately 1800 cm³ of fecal matter containing 7082 seeds. The fecal collections from the 3 early wet season locations were deposited by tapirs in moist areas that were not flooded at the time of collection. Thirteen of the 20 mid-wet season samples were intact boluses collected from a flooded drainage and 7 were collected immediately adjacent to this drainage on dry soil.

In total, the feces contained 8166 seeds of at least 16 plant species. The size range of seeds encountered was approximately 1–30 mm in length, with 7864 small seeds, 193 medium seeds and 109 large seeds. The majority of the seeds (96%) and species (50%) were small-seeded. As much as 83% of all seeds came from the canopy tree *Laetia procera* (Flacourtiaceae); these 2–3 mm sized seeds were present in each of the 20 fecal samples collected in August. The next most abundant (group of) species was *Ficus*, which made up 11% of the seeds. Of all small seeds (<5 mm), the majority (69%) contained an intact endosperm.

DISCUSSION

Tapirs are important dispersers of large-seeded tree species (Bodmer 1991; Fragoso 1997; Quiroga-Castro & Roldan 2001; Fragoso *et al.* 2003; Giombini *et al.* 2009; O’Farrill *et al.* 2012). Previous studies reported that *T. terrestris* feces contain large numbers of small seeds (e.g. Fragoso & Huffman 2000; Tófoli 2006; Talamoni & Assis 2009; Zorzi 2009; Tobler *et al.* 2010). Our study provides similar data for *T. bairdii* and suggests that a large

proportion of small seeds that pass through the digestive tract of *T. bairdii* are viable. Feces of the Central American tapir on Barro Colorado Island contained large numbers of small seeds, which by far outnumbered large seeds in the samples. The majority of these appeared viable in cut tests.

The most abundant seed in the fecal samples we collected was *L. procera*, an uncommon species in the old-growth forest of BCI, which produces 15–20 mm large berrylike capsules containing 10–20 seeds, in Aug–Sep (Croat 1978). These berrylike fruits are rich in crude fat and energy content (Castellanos & Chanin 1996) and are known to be eaten by a variety of birds and primates that collect the fruits directly from the trees (Snow 1971; Van Roosmalen *et al.* 1988; Castellanos & Chanin 1996; Wehncke *et al.* 2003). The tapirs producing the feces of this study must have fed on *L. procera* fruits that had fallen to the forest floor and deposited the seeds at latrines 79% intact.

As our study covered just 4 different latrines and possibly few individual tapirs, the abundances of various plant species we recorded in feces might not be representative of an unbiased sample. For example, *L. procera* appears to be overrepresented. This study did not address the relationship between abundance of seeds found in feces and the availability of the same species at the time of fecal sample collection. Year-round sampling in combination with large-scale inventory and monitoring of tree fruiting phenology would allow us to assess whether tapirs feed selectively, or rather on whatever is available.

Our results suggest that tapirs might act as dispersers not only of large-seeded plant species, as documented by previous studies (Bodmer 1991; Rodrigues *et al.* 1993; Fragoso 1997; Fragoso & Huffman 2000; Quiroga-Castro & Roldan 2001; Giombini *et al.* 2009; Tobler *et al.* 2010; O’Farrill *et al.* 2012), but also of plant species with relatively small seeds that are not presently considered ‘tapir-dispersed’. To assess this role with more certainty, it would be worthwhile investigating the occurrence and viability rates of small seeds (<5 mm) in tapir dung at other locations and in all seasons. In addition, although cut tests provide acceptable estimates of viability (Terry *et al.* 2003; as in Fragoso & Huffman 2000, Daws *et al.* 2002 and Strong & Fragoso 2006), germination tests are recommended to determine viability with greater certainty.

Ultimately, the effectiveness of tapirs as dispersers of small seeds will also depend on the fate of the vi-

Table 1 Seeds found within 72 tapir feces of Central American tapir (*Tapirus bairdii*) on Barro Colorado Island, Panama

Species (family)	Life form	Seed size (length), size class [†]	Fresh mass (g) [‡]	Number	Number per feces (mean \pm SD)	Number intact	Estimated viability (%)	Frequency (% of feces) [§]
<i>Ficus</i> sp. (Moraceae)	Tree/midstory	1 mm, S	NA	882	12.3 (\pm 21.5)	75	9	59.7
<i>Casearia</i> sp. (Flacourtiaceae)	Tree	1 mm, S	NA	6	0.08 (\pm 0.3)	2	33	6.9
Unknown sp ₁		1 mm, S	NA	23	0.3 (\pm 0.8)	1	4	13.9
Unknown sp ₂		1 mm, S	NA	100	1.4 (\pm 3.3)	3	3	19.4
Unknown sp ₃		1 mm, S	NA	4	0.05 (\pm 0.2)	2	50	5.6
Unknown sp ₄		2 mm, S	NA	32	0.4 (\pm 1.0)	25	78	25.0
Unknown sp ₅		2 mm, S	NA	22	0.3 (\pm 0.7)	4	18	20.8
<i>Laetia procera</i> (Flacourtiaceae)	Tree	2–3 mm, S	0.007	6795	94.4 (\pm 195.2)	5336	79	27.8
<i>Psychotria</i> sp. (Rubiaceae)	Shrub	5 mm, M	0.01–0.03	176	2.4 (\pm 5.4)	0	0	25.0
<i>Meibomia axillaris</i> var. <i>Acutifolia</i> (Fabaceae)	Herbaceous	5 mm, M	NA	4	0.05 (\pm 0.2)	0	0	5.6
Unknown sp ₆		5 mm, M	NA	1	0.01 (\pm 0.1)	0	0	1.4
Fabacea sp.	Shrub/tree	8 mm, M	NA	6	0.08 (\pm 0.3)	6	100	8.3
<i>Cayaponia</i> sp. (Cucurbitaceae)		8 mm, M	NA	3	0.04 (\pm 0.2)	0	0	4.2
Unknown sp ₇		15 mm, M	NA	3	0.04 (\pm 0.2)	2	67	4.2
<i>Spondias mombin</i> (Anacardiaceae)	Tree	20–23 mm, L	1.9	108	1.5 (\pm 2.7)	102	94	27.8
<i>Astrocaryum standleyanum</i> (Arecaceae)	Tree (palm)	30 mm, L	9.7	1	0.01 (\pm 0.1)	1	100	1.4
Total				8166	113.4 (\pm 195.2)	5559	68.1	
Mean								39.7 (\pm 40.4) [¶]

[†]Seed sizes are approximate. Size class (length): S < 5 mm, M = 5–20 mm, L > 20 mm. [‡]Fresh mass of *Laetia procera* obtained from Guariguata (2000); all other fresh mass records provided by J. Wright (pers. comm.). [§]Frequency = percentage of total fecal samples that contained the species. [¶]Mean estimated viability (%) across species. NA, not available.

able seeds in the feces in relation to the habitat. Tapirs are known to deposit feces in both dry and flooded sites (Fragoso & Huffman 2000; Tobler *et al.* 2010). For flooded sites, tapir-mediated dispersal might be especially effective for plant species associated with wet habitats (Salas & Fuller 1996), to which tapirs may provide directed dispersal (Wenny 2001). Dispersal by tapirs might also be effective for other small-seeded plant species when combined with secondary seed dispersal by dung beetles that move and bury dung in the soil (e.g. seeds in howler monkey feces [Estrada & Coates-Estrada 1991]).

ACKNOWLEDGMENTS

We thank the Smithsonian Tropical Research Institute for hosting this study; Jim Dalling for use of his laboratory; Osvaldo Calderón and Andres Hernandez for seed identification; Eloisa Lasso De Paulis for assistance in the field; Ricardo Moreno for assistance in the field and help with herbarium seed comparison; José M. V. Fragoso and Andrew Taylor for advice; Joe Wright for seed mass data; Roland Kays for current BCI tapir population information; and José M. V. Fragoso, Aaron Shiels, Laura Capece, Michael Szperka for comments on previous versions of the manuscript. Patrick A. Jansen acknowledges financial support from the Netherlands Foundation for Scientific Research.

REFERENCES

- Bodmer RE (1991). Strategies of seed dispersal and seed predation in Amazonian ungulates. *Biotropica* **23**, 255–61.
- Castellanos HG, Chanin P (1996). Seasonal differences in food choice and patch preference of long-haired spider monkeys (*Ateles belebuth*). In: Norconk MA, Rosenberger AL, Garber PA, eds. *Adaptive Radiations of Neotropical Primates*. Plenum Press, New York, pp. 451–66.
- Croat TB (1978). *Flora of Barro Colorado Island*. Stanford University Press, Stanford, CA.
- Daws MI, Burslem DFRP, Crabtree LM, Kirkman P, Mullins CE, Dalling JW (2002). Differences in seed germination responses may promote coexistence of 4 sympatric piper species. *Functional Ecology* **16**, 258–67.
- Eisenberg JF (1980). The density and biomass of tropical mammals. In: Soule ME, Wilcox BA, eds. *Conservation Biology: An Evolutionary-Ecological Perspective*. Sinauer Associates, Sunderland, MA, pp. 35–55.
- Eisenberg JF, Thorington RW (1973). A preliminary analysis of a Neotropical mammal fauna. *Biotropica* **5**, 150–61.
- Enders RK (1935). Mammalian life histories from Barro Colorado Island, Panama. *Bulletin of the Museum of Comparative Zoology* **78**, 385–502.
- Enders RK (1939). Changes observed in the mammal fauna of Barro Colorado Island, 1929–37. *Ecology* **20**, 104–6.
- Estrada A, Coates-Estrada R (1991). Howler monkeys (*Alouatta palliata*), dung beetles (Scarabaeidae) and seed dispersal: ecological interactions in the tropical rain forest of Los Tuxtlas, Mexico. *Journal of Tropical Ecology* **7**, 459–74.
- Flesher K (1999). Preliminary notes on the conservation status of Bair's tapir *Tapirus bairdii* in northeastern Honduras. *Oryx* **33**, 294–300.
- Foerster CR, Vaughan C (2002). Home range, habitat use and activity of Baird's tapir in Costa Rica. *Biotropica* **34**, 423–37.
- Forget PM, Sabatier D (1997). Dynamics of the seedling shadow of a frugivore-dispersed tree species in French Guiana. *Journal of Tropical Ecology* **13**, 767–73.
- Fragoso JMV (1997). Tapir-generated seed shadows: scale-dependent patchiness in the Amazon rain forest. *Journal of Ecology* **85**, 519–29.
- Fragoso JMV, Huffman JM (2000). Seed-dispersal and seedling recruitment patterns by the last Neotropical megafaunal element in Amazonia, the tapir. *Journal of Tropical Ecology* **16**, 368–85.
- Fragoso JMV, Silviu KM, Correa JA (2003). Long-distance seed dispersal by tapirs increased seed survival and aggregates tropical trees. *Ecology* **84**, 1996–2006.
- Giombini MI, Bravo SP, Martinez MF (2009). Seed dispersal of the palm *Syagrus romanzoffiana* by tapirs in the semi deciduous Atlantic forest of Argentina. *Biotropica* **41**, 408–13.
- Glanz WE (1996). The terrestrial mammal fauna of Barro Colorado Island: censuses and long term changes. In: Leigh E, Rand S, Windsor D, eds. *The Ecology of a Tropical Forest: Seasonal Rhythms and Long term Changes*, 2nd edn. Smithsonian Tropical Research Institute, Washington, DC, pp. 455–68.
- Guariguata MR (2000). Seed and seedling ecology of tree species in Neotropical secondary forests: management implications. *Ecological Applications* **10**, 145–54.

- Henry O, Feer F, Sabatier D (2000). Diet of the lowland tapir (*Tapirus terrestris* L.) in French Guiana. *Biotropica* **32**, 364–8.
- IUCN (2012). *The IUCN Red List of Threatened Species*. Version 2012.1. [Cited 18 Jul 2012.] Available from URL: www.iucnredlist.org
- Janzen DH (1981). Digestive seed predation by a Costa Rican Baird's tapir. *Biotropica* **13**, 59–63.
- Janzen DH (1982). Seeds in tapir dung in Santa Rosa National Park, Costa Rica. *Brenesia* **19/20**, 129–35.
- Janzen DH (1984). Dispersal of small seeds by big herbivores: foliage is the fruit. *The American Naturalist* **123**, 339–53.
- Jordano P, Schupp EW (2000). Seed disperser effectiveness: the quantity component and patterns of seed rain for *Prunus mahaleb*. *Ecological Monographs* **70**, 591–615.
- Leigh E, Rand S, Windsor D (1996). *The Ecology of a Tropical Forest: Seasonal Rhythms and Long term Changes*, 2nd edn. Smithsonian Tropical Research Institute, Washington, DC.
- Naranjo EJ (1995). Hábitos de alimentación del tapir (*Tapirus bairdii*) en un bosque tropical húmedo de Costa Rica. *Vida Silvestre Neotropical* **4**, 32–7.
- Naranjo EJ (2009). Ecology and conservation of Baird's tapir in Mexico. *Tropical Conservation Science* **2**, 140–58.
- O'Farrill G, Calmé S, Gonzalez A (2006). *Manilkara zapota*: a new record of a species dispersed by tapirs. *Tapir Conservation* **15**, 32–5.
- O'Farrill G, Calmé S, Sengupta R, Gonzalez A (2012). Effective dispersal of large seeds by Baird's tapir: a large-scale field experiment. *Journal of Tropical Ecology* **28**, 119–22.
- Olmos F (1997). Tapirs as seed dispersers and predators. In: Brooks DM, Bodmer RE, Matola S, eds. *Tapirs: Status Survey and Conservation Action Plan*. IUCN, Gland, Switzerland, pp. 3–9.
- Quiroga-Castro VD, Roldan AI (2001). The fate of *Attalea phalerata* (Palmae) seeds dispersed to a tapir latrine. *Biotropica* **33**, 472–7.
- Rodrigues M, Olmos F, Galetti M (1993). Seed dispersal by tapir in southeastern Brazil. *Mammalia* **57**, 460–61.
- Salas LA, Fuller TK (1996). Diet of the lowland tapir (*Tapirus terrestris* L.) in the Tabaro River valley, southern Venezuela. *Canadian Journal of Zoology* **74**, 1444–51.
- Schupp EW (1993). Quantity, quality and the effectiveness of seed dispersal by animals. *Vegetatio* **107/108**, 15–29.
- Snow BK, Snow DW (1971). The feeding ecology of tanagers and honeycreepers in Trinidad. *The Auk* **88**, 291–322.
- Strong JN, Fragoso JMV (2006). Seed dispersal by *Geochelone carbonaria* and *Geochelone denticulata* in northwestern Brazil. *Biotropica* **38**, 683–6.
- Talamoni SA, Assis MAC (2009). Feeding habit of the Brazilian tapir, *Tapirus terrestris* (Perrissodactyla: Tapiridae) in a vegetation transition zone in southeastern Brazil. *Zoologia Curitiba Impresso* **26**, 251–4.
- Terry J, Probert RJ, Linington SH (2003). Processing and maintenance of the Millennium Seed Bank collections. In: Smith RD, Dickie JB, Linington SH, Pritchard HW, Probert RJ, eds. *Seed Conservation. Turning Science into Practice*. Royal Botanic Gardens, Kew, pp. 307–25.
- Terwilliger V (1978). Natural history of Baird's tapir on Barro Colorado Island, Panama Canal Zone. *Biotropica* **10**, 211–20.
- Tobler MW (2002). Habitat use and diet of Baird's tapirs (*Tapirus bairdii*) in a montane cloud forest of Cordillera de Talamanca, Costa Rica. *Biotropica* **34**, 468–74.
- Tobler MW, Janovec JP, Cornejo F (2010). Frugivory and seed dispersal by the lowland tapir *Tapirus terrestris* in the Peruvian Amazon. *Biotropica* **42**, 215–22.
- Tófoli CF (2006). Frugivoria e dispersão de sementes por *Tapirus terrestris* (Linnaeus, 1758) na paisagem fragmentada do Pontal do Paranapanema, São Paulo (MS thesis). University of São Paulo, São Paulo, Brazil.
- Van Roosmalen MGM, Mittermeier RA, Fleagle JG (1988). Diet of the northern bearded saki (*Chiropotes satanas chiropotes*): a Neotropical seed predator. *American Journal of Primatology* **14**, 11–35.
- Wehncke EV, Hubbell SP, Foster RB, Dalling JW (2003). Seed dispersal patterns produced by white-faced monkeys: implications for the dispersal limitation of Neotropical tree species. *Journal of Ecology* **91**, 677–85.
- Wenny DG (2001). Advantages of dispersal: a re-evaluation of directed dispersal. *Evolutionary Ecology Research* **3**, 51–74.
- Wright JS, Gompper ME, DeLeon B (1994). Are large predators keystone species in Neotropical forests? The evidence from Barro Colorado Island. *Oikos* **71**, 279–94.
- Zorzi BT (2009). Frugivoria por *Tapirus terrestris* em três regiões do Pantanal, Brasil (MS thesis). Federal University of Mato Grosso do Sul, Campo Grande, Brazil.